arXiv:1303.4548v1 [math.PR] 19 Mar 2013

BASIC PROPERTIES OF CRITICAL LOGNORMAL MULTIPLICATIVE CHAOS

JULIEN BARRAL, ANTTI KUPIAINEN[1,2], MIIKA NIKULA[1], EERO SAKSMAN[1],
AND CHRISTIAN WEBB[1]

Abstract. We study one-dimensional exact scaling lognormal multiplicative chaos measures at criticality. Our main results are the determination of the exact asymptotics of the right tail of the distribution of the total mass of the measure, and an almost sure upper bound for the modulus of continuity of the cumulative distribution function of the measure. We also find an almost sure lower bound for the increments of the measure almost everywhere with respect to the measure itself, strong enough to show that the measure is supported on a set of Hausdorff dimension 0.

1. Introduction

Multiplicative chaos is a theory developed by Kahane in the eighties [25, 23, 26]. It deals with multiplicative processes generating martingales, which take values in the cone of non-negative Radon measures on σ-compact metric spaces. This theory is based on the lognormal multiplicative chaos proposed by Mandelbrot to model turbulence [31], as well as the works previously achieved by Kahane and Peyrière [28] on the simplified model of multiplicative cascades on trees still proposed by Mandelbrot [32, 33], namely the so-called Mandelbrot cascades, which assume no log-normality property. The study of random measures generated by such multiplicative processes also originates from random covering and percolation theory questions (see [24, 23, 26, 27, 22, 5]). When statistically self-similar, as it is the case for limits of Mandelbrot cascades, these measures provide nice illustrations of the so-called multifractal formalism, as well as models in the study of intermittent phenomena beyond turbulence, like the distribution of rare minerals in earth [34] or stock exchange fluctuations in finance [35]. Examples of such measures on $\mathbb{R}^d$ possessing continuous (rather than only discrete for limits of Mandelbrot cascades) scaling properties are some of the Gaussian multiplicative chaos built by Kahane in [25] or the Lévy multiplicative chaos built by Fan in [21], the compound Poisson cascades built by Barral and Mandelbrot [9] and their generalization in the so-called infinitely divisible cascades by Bacry and Muzy in [4].

Kahane's lognormal multiplicative chaos has been recently revisited and completed in several directions [39, 40, 2]. Also, it is now a central tool in two-dimensional quantum gravity theory since it provides, through the

---

Date: March 20, 2013.

2010 *Mathematics Subject Classification.* Primary 60G57; Secondary 60G18,83C45.

Key words and phrases. Multiplicative chaos, critical temperature.

[1]Supported by the [1]Academy of Finland and [2]ERC

exponential of the Gaussian free field, the random measures used to obtain the first rigorous results in direction to the so-called KPZ formula in works by Duplantier and Sheffield [16, 17], as well as Rhodes and Vargas [38] (see also Benjamini and Schramm [11] for a 1-dimensional version in the framework of Mandelbrot multiplicative cascades on $[0, 1]$). Nondegenerate limits of lognormal multiplicative chaos associated with the exponential of the Gaussian free field on the circle have also been used successfully by Astala, Jones, Kupiainen and Saksman in [3] to build random planar curves by conformal welding. The families of Gaussian multiplicative chaos considered in these questions are naturally parametrized by a continuous parameter $\beta \in [0, \beta_c)$ which, in the quantum gravity is in bijection with the so-called central charge, in random energy models corresponds to the inverse of a temperature, in turbulence is a measure of the intermittence, and is a decreasing function of the Hausdorff dimension of the associated measure in the Euclidean geometry. At the critical temperature, and below it, the limit μ_β of the martingale $\mu_{\beta,t}$ provided by the associated multiplicative process vanishes almost surely. For $\beta > \beta_c$, it is nevertheless possible to give a sense to the corresponding dual KPZ formula [15, 7] by considering measures essentially by subordinating a suitable nondegenerate Gaussian multiplicative chaos to some stable Lévy subordinators; this yields an atomic measure. At the critical value β_c, one needs new results in multiplicative chaos theory. They were obtained during the last months by Duplantier, Rhodes, Sheffield and Vargas in [18, 19], inspired by results recently achieved by Aïdékon and Shi in the context of the martingales in the branching random walk [1]. Thus, it is possible to get a nontrivial positive measure at the critical temperature as the limit of the signed measures $-\frac{\mathrm{d}\mu_{\beta,t}}{\mathrm{d}\beta}_{|\beta=\beta_c}$ as $t \to \infty$. Moreover, this measure is continuous. We also mention that like in the context of martingales in the branching random walks [1], the critical measure can be obtained as limit in probability of $\mu_{\beta_c,t}$ properly normalized [19], while for $\beta > \beta_c$ such normalization results are still unknown though suspected to be analogous to those known in the branching random walk and random energy models frameworks [41, 30, 8]; in this sense, it is reasonable to believe that for $\beta > \beta_c$, the location of the atoms of the weak limit of $\mu_{\beta,t}$ properly renormalized is in an essential way governed by the critical measure distribution, and consequently any information about the critical measure might be useful for the low temperature measures.

This paper is dedicated to the study of some properties of such critical lognormal multiplicative chaos measure. We concentrate on the exactly scale-invariant one dimensional construction. Our main results are the determination of the asymptotic behavior of the tail of the distribution of the total mass of the measure, a bound for the modulus of continuity of the measure for which the previous tail asymptotic behavior is crucial, and an estimate from below of the measure increments almost everywhere with respect to the measure, which completes the estimation provided by the modulus of continuity and goes beyond the simple fact that the measure has Hausdorff dimension 0, see Theorems 1, 2 and 3 below.

1.1. **Definitions and notation.** In this section we fix notation and give the precise definitions of the objects studied in this paper. Formally, the one-dimensional lognormal multiplicative chaos measures μ_β are random measures given by

$$\mu_\beta(\,\mathrm{d}x) = e^{\beta X(x) - \frac{\beta^2}{2}\mathbb{E}X(x)^2}\,\mathrm{d}x, \tag{1}$$

where $(X(x))_{x\in\mathbb{R}}$ is a logarithmically correlated centered Gaussian field, i.e. a centered Gaussian process with

$$\mathbb{E}X(x)X(y) \sim \log\frac{1}{|x-y|} \quad \text{as } |x-y| \to 0.$$

However, the logarithmic singularity of the correlation kernel implies that the realizations of X are not smooth enough to be functions, but must instead be defined as random distributions. To overcome this major technical obstacle, in Kahane's theory of multiplicative chaos one gives a rigorous meaning to the expression (1) by considering nonsingular approximations X_t to the field X, defining the measures $\mu_{\beta,t}$ corresponding to these regularizations and then taking the weak limit of the measures $\mu_{\beta,t}$ as the approximation parameter is taken to infinity. In this way one completely avoids the problem of defining the exponential of a distribution.

We mainly concentrate on the *exactly scale invariant* construction. This scaling property, to be defined below, is central to the proof of Theorem 1. The one-dimensional exactly scale invariant construction is most easily understood through the following geometric construction, originally due to Bacry and Muzy [4].

Let λ denote the hyperbolic area measure on the upper half-plane, i.e.

$$\lambda(A) = \int_A \frac{\mathrm{d}x\,\mathrm{d}y}{y^2} \quad \text{for all } A \subset \mathbb{R}\times\mathbb{R}^+.$$

For $x\in\mathbb{R}$ and $t\in\mathbb{R}_+$, let $\mathcal{C}_t(x)$ denote the set

$$\mathcal{C}_t(x) = \left\{(x',y') \,\middle|\, y' > \max(2|x'-x|, e^{-t}), |x'-x| < \frac{1}{2}\right\}$$

and for a compact interval $I \subset \mathbb{R}$ of length less than or equal to 1, denote

$$\mathcal{C}_t(I) = \bigcap_{x\in I} \mathcal{C}_t(x).$$

Note that for $t \geq \log 1/|I|$ we have $\mathcal{C}_t(I) = \mathcal{C}_{\log 1/|I|}(I)$. Next, let W denote the white noise on $\mathbb{R}\times\mathbb{R}^+$ with control measure λ. We consider W a random real function on the Borel sets of $\mathbb{R}\times\mathbb{R}^+$ with finite λ-measure characterized by the following properties: for all disjoint Borel sets $A, B \subset \mathbb{R}\times\mathbb{R}^+$ such that $\lambda(A), \lambda(B) < \infty$,

1. $W(A)$ is a centered Gaussian random variable with variance $\lambda(A)$,
2. the random variables $W(A)$ and $W(B)$ are independent, and
3. almost surely we have $W(A\cup B) = W(A) + W(B)$.

Define

$$X_t(x) = W(\mathcal{C}_t(x)) \quad \text{for all } x\in\mathbb{R}, t\in[0,\infty).$$

For a fixed $t > 0$, the covariance structure of the process $(X_t(x))_{x\in\mathbb{R}}$ can be computed to be

$$\mathbb{E}X_t(x)X_t(y) = \begin{cases} t+1-e^t|x-y|, & |x-y| < e^{-t} \\ \log\frac{1}{|x-y|}, & e^{-t} \le |x-y| \le 1 \\ 0, & 1 < |x-y| \end{cases} .$$

For any interval $I \subset \mathbb{R}$ of length less than or equal to 1 and $x \in I$ we denote

$$X_t(I) = W(\mathcal{C}_t(I)) \quad \text{and} \quad X_t^I(x) = W(\mathcal{C}_t(x) \setminus \mathcal{C}_t(I))$$

to obtain the decomposition $X_t(x) = X_t(I) + X_t^I(x)$, where $X_t(I)$ is independent of the process $(X_t^I(x))_{x\in I}$. Since $\mathcal{C}_t(I) = \mathcal{C}_{\log 1/|I|}(I)$ for $t \ge \log 1/|I|$, we denote $X(I) := X_{\log 1/|I|}(I)$. Owing to the geometry of the construction, the field $(X_t(x))$ satisfies the following scale invariance property: for all intervals $I \subset \mathbb{R}$ and $e^{-t} < |I| < 1$ we have

(2)
$$(X_t(x))_{x\in I} = \left(X_t(I) + X_t^I(x)\right)_{x\in I} \overset{d}{=} \Big(X_t(I) + X'_{t-\log|I|}(x/|I|)\Big)_{x\in[0,1]},$$

where X' is an independent realization of the field X. For reader's convenience we give the geometric explanation for this scaling property in the appendix.

For $\beta \in (0, \sqrt{2})$ we construct the measures $\mu_{\beta,t}$ on the unit interval by setting

$$\mu_{\beta,t}(I) = \int_I e^{\beta X_t(x) - \frac{\beta^2}{2}\mathbb{E}X_t(x)^2}\,\mathrm{d}x \tag{3}$$

for all intervals $I \subset [0,1]$. This construction fits into the framework of Kahane's theory of multiplicative chaos [23], which implies that almost surely the limit $\mu_\beta = \lim_{t\to\infty}\mu_{\beta,t}$ exists in the sense of weak convergence of measures and that the limit measure satisfies $\mu_\beta(I) > 0$ for all intervals $I \subset [0,1]$. The scaling property (2) implies that the measures μ_β are exactly scale invariant, especially that

$$\mu_\beta(I) \overset{d}{=} |I| e^{\beta X(I) - \frac{\beta^2}{2}\mathbb{E}X(I)^2}\mu'_\beta([0,1]) \quad \text{for all intervals } I \subset [0,1], \tag{4}$$

where μ'_β is an independent realization of μ_β and $X(I)$, defined as above, is a centered Gaussian random variable of variance $\log\frac{1}{|I|}$.

Kahane's work also implies that the corresponding construction for $\beta \ge \sqrt{2}$ results in degenerate limit measures, i.e. the limit measure will be almost surely null. However, the exact scaling relation above makes sense for all $\beta > 0$. It has recently been shown by Duplantier, Rhodes, Sheffield and Vargas [18] that by defining

(5)
$$\begin{aligned} \mu_{\sqrt{2},t}(I) &= -\left.\frac{\mathrm{d}}{\mathrm{d}\beta}\right|_{\beta=\sqrt{2}} \mu_{\beta,t}(I) \\ &= \int_I (\sqrt{2}(t+1) - X_t(x))e^{\sqrt{2}X_t(x) - \mathbb{E}X_t(x)^2}\,\mathrm{d}x \quad \text{for all intervals } I \subset [0,1] \end{aligned}$$

one obtains a nondegenerate almost sure weak limit $\mu_{\sqrt{2}} = \lim_{t\to\infty}\mu_{\sqrt{2},t}$ for which $\mu_{\sqrt{2}}(I) > 0$ almost surely for all intervals $I \subset [0,1]$. As in the case of branching random walks (or equivalently, multiplicative cascades), this derivative turns out to be the correct replacement for the measures (3) in the case $\beta = \sqrt{2}$, at the very least in the sense that $\mu_{\sqrt{2}}$ is nontrivial and turns out to satisfy the exact scaling property: as detailed in the appendix, we have especially

$$\mu_{\sqrt{2}}(I) \overset{d}{=} |I| e^{\sqrt{2}X(I) - \mathbb{E}X(I)^2} \mu'_{\sqrt{2}}([0,1]) \quad \text{for all intervals } I \subset [0,1]. \tag{6}$$

In defining the lognormal multiplicative chaos measure for the critical parameter value $\beta = \sqrt{2}$, the peculiar normalizing factor $(\sqrt{2}(t+1) - X_t(x))$ may also be replaced by a normalization that is deterministic and also independent of x. Inspired by the arguments of Aïdékon and Shi [1] in the case of branching random walks, Duplantier, Rhodes, Sheffield and Vargas [19] recently proved that there exists a deterministic constant $c > 0$ such that for every interval $I \subset [0,1]$ one has

$$\sqrt{t} \int_I e^{\sqrt{2}X_t(x) - \mathbb{E}X_t(x)^2}\, \mathrm{d}x \to c\, \mu_{\sqrt{2}}(I) \quad \text{in probability as } t \to \infty. \tag{7}$$

Before moving on to the statements of our results on the fine properties of $\mu_{\sqrt{2}}$ we make a final comment on the scale invariance properties of multiplicative chaos measures. In [18] and [19] the authors deal primarily with a slightly different construction, the $\star$-*scale invariant* lognormal multiplicative chaos measures. In the terms of the geometric construction presented here, a $\star$-scale invariant random measure is obtained by replacing the field $(X_t(x))$ in (3), (5) or (7) by the field $(X_t(x) - X_0(x))$. Since we will make use of this construction in the proof of Theorem 2, we have included details on $\star$-scale invariance in the appendix. However, as also noted in the papers themselves, the proofs of the convergence results in [18] and [19] are insensitive to these differences.

1.2. **Main results.**

We will make use of the following result of Duplantier, Rhodes, Sheffield and Vargas [19], which is a corollary of the deterministic normalization (7).

Theorem A. *For all* $h \in (0,1)$, $\mathbb{E}(\mu_{\sqrt{2}}([0,1]))^h < \infty$.

The first of our main theorems is a strengthening of this result, and analogous to the theorem of Buraczewski [13] on the fixed points of the smoothing transform.

Theorem 1. *The tail probabilities of* $\mu_{\sqrt{2}}$ *have the asymptotic behavior*

$$\mathbb{P}\left(\mu_{\sqrt{2}}([0,1]) > \lambda\right) \sim c_1/\lambda \quad \text{as } \lambda \to \infty,$$

where the constant is given explicitly by

$$c_1 = \frac{2}{\log 2} \mathbb{E}\mu_{\sqrt{2}}([0,1/2]) \log\left(1 + \frac{\mu_{\sqrt{2}}([1/2,1])}{\mu_{\sqrt{2}}([0,1/2])}\right) < \infty.$$

This theorem allows one to get detailed information on the geometric properties of the measure $\mu_{\sqrt{2}}$. The following result is analogous to our earlier result [10] on multiplicative cascades.

Theorem 2. *For any interval* $I \subset [0,1]$ *and* $\gamma < \frac{1}{2}$

$$\mu_{\sqrt{2}}(I) \le C(\omega)\left(\log(1+|I|^{-1})\right)^{-\gamma}, \tag{8}$$

where $C(\omega) > 0$ *is an almost surely finite random constant.*

The proof of this theorem is inspired by the earlier result, but as the correlations of the field X in the construction of $\mu_{\sqrt{2}}$ are much more intricate than in the branching random walk underlying the cascade measures, more involved arguments are needed.

We also get a bound on the appropriate Hausdorff gauge function to measure the size of the smallest Borel sets fully supporting $\mu_{\sqrt{2}}$. We have the following result.

Theorem 3. *Denote* $f_\alpha(n) = \exp\left(-\sqrt{6\log 2}\sqrt{n\left(\log n + \alpha \log\log n\right)}\right)$ *for* $\alpha > \frac{1}{3}$. *Almost surely,*

$$\mu_{\sqrt{2}}\big(\{x : \mu_{\sqrt{2}}(I_n(x)) \ge f_\alpha(n) \textit{ for all but finitely many } n\}\big) = \mu_{\sqrt{2}}([0,1]).$$

The proof uses large deviations estimates exploiting both the exact scaling property of $\mu_{\sqrt{2}}$ and the tail probabilities given by Theorem 1. This theorem implies the weaker claim that almost surely there exists a set of full $\mu_{\sqrt{2}}$-measure that has Hausdorff dimension 0, a fact that we state as Corollary 17.

For the log-normal critical Mandelbrot measure μ on trees, we establish in [10] that $\mu\big(\{x : \mu(I_n(x)) \le \psi(n) \text{ for all but finitely many } n\}\big) = \mu([0,1])$, for all functions $\psi(n) = n^{-k}$, $k \ge 1$. In particular the modulus of continuity (shown to be optimal) does not capture the measure increments behavior μ-almost everywhere. The proof exploits fine information about the partition functions associated with μ at low temperatures. This information strongly depends on the renormalization theory for the low temperature measures $\mu_{\beta,n}$. As mentioned above, it is believed that they have their analogues in the present setting, but establishing them remains a challenge, as well as proving the optimality of the bound provided by Theorem 2. Anyway, if these properties hold for $\mu_{\sqrt{2}}$, as for discrete cascades on trees it remains to determine whether they are sharp or not. At least they draw the contours of the asymptotic behavior of these measures increments almost everywhere with respect to themselves, which must differ from the law of the iterated logarithm observed in the high temperature case (see [6], and [29] for measures on the boundary of Galton-Watson trees).

2. TAIL PROBABILITIES

The proof of Theorem 1 follows the same idea as the earlier closely related results of Durrett and Liggett [14], Guivarc'h [20], Liu [29], Buraczewski [13, 12] and Barral and Jin [6]: one uses the smoothing transform (or in the case of multiplicative chaos, a similar distributional equation with more dependencies) to derive a Poisson equation satisfied by the quantity one is interested in, and then analyzes the behavior of the solutions of the Poisson equation at infinity. A key point in the derivations of the Poisson equations in all these proofs is the use of an alternate probability measure (the Peyrière probability), the idea of which goes back to the seminal paper of Kahane and Peyrière [28].

In this section we denote $\mu := \mu_{\sqrt{2}}$ and $Y := \mu([0,1])$. The variable Y may be written as the fixed point of a "non-independent smoothing transform" as follows:

$$\begin{aligned} Y = \mu([0,1]) &= \mu([0,1/2]) + \mu([1/2,1]) \\ &=: \frac{1}{2}e^{\sqrt{2}X([0,1/2]) - \mathbb{E}X([0,1/2])^2}Y_0 + \frac{1}{2}e^{\sqrt{2}X([1/2,1]) - \mathbb{E}X([1/2,1])^2}Y_1 \\ &=: W_0Y_0 + W_1Y_1. \qquad (9) \end{aligned}$$

By the exact scale invariance property (6) of μ we have $Y_0 \perp W_0$, $Y_1 \perp W_1$ and $Y_0 \overset{d}{=} Y \overset{d}{=} Y_1$. Note, however, that Y_0 is not independent of either Y_1 or W_1.

We then define the version of Peyrière probability that is most convenient for our needs.

Definition 4. *Let $(\Omega, \mathcal{F}, \mathbb{P})$ denote the probability space on which μ is defined and define a probability space $(\Omega \times \{0,1\}, \mathcal{F} \times \sigma(\{0,1\}), \mathbb{Q})$ by setting*

$$\mathbb{E}_{\mathbb{Q}} f(\omega, j) = \mathbb{E}\left(W_0(\omega)f(\omega,0) + W_1(\omega)f(\omega,1)\right)$$

for all bounded $\mathcal{F} \times \sigma(\{0,1\})$-measurable functions $f : \Omega \times \{0,1\}$. Define the random variables $\widetilde{Y}$, $\widetilde{W}$ and $\widetilde{B}$ on this probability space by setting

$$\widetilde{Y}(\omega,j) = Y_j(\omega), \quad \widetilde{W}(\omega,j) = W_j(\omega) \text{ and } \widetilde{B}(\omega,j) = \begin{cases} W_1(\omega)Y_1(\omega), & j = 0 \\ W_0(\omega)Y_0(\omega), & j = 1 \end{cases}.$$

We state the essential properties of these variables as the following lemma.

Lemma 5. (1) *$\widetilde{W}$ and $\widetilde{Y}$ are independent.*

(2) *$\widetilde{Y}$ (under $\mathbb{Q}$) has the same law as Y (under $\mathbb{P}$).*

(3) *$-\log \widetilde{W}$ is a centered Gaussian with variance $2\log 2$.*

Proof. Let $f, g : \mathbb{R} \mapsto \mathbb{R}$ be bounded and continuous. By direct computation and the independences $W_0 \perp Y_0$ and $W_1 \perp Y_1$,

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}} f(\widetilde{W})g(\widetilde{Y}) &= \mathbb{E}\big(W_0f(W_0)g(Y_0) + W_1f(W_1)g(Y_1)\big) \\ &= 2\big(\mathbb{E}W_0f(W_0)\big)\big(\mathbb{E}g(Y_0)\big). \end{aligned}$$

By taking $g \equiv 1$ we see that $\mathbb{E}_{\mathbb{Q}} f(\widetilde{W}) = 2\mathbb{E}W_0f(W_0)$, and taking $f \equiv 1$ yields $\mathbb{E}_{\mathbb{Q}} g(\widetilde{Y}) = \mathbb{E}g(Y_0) = \mathbb{E}g(Y)$. Thus (2) holds and moreover

$$\mathbb{E}_{\mathbb{Q}} f(\widetilde{W})g(\widetilde{Y}) = \mathbb{E}_{\mathbb{Q}} f(\widetilde{W})\mathbb{E}_{\mathbb{Q}} g(\widetilde{Y}),$$

which means that $\widetilde{W}$ and $\widetilde{Y}$ are independent as claimed in (1).

The law of $-\log \widetilde{W}$ is easy to identify by computing the moment generating function. Since $W_0 \overset{d}{=} W_1 \overset{d}{=} \frac{1}{2}e^{\sqrt{2\log 2}N - \log 2}$, where N is a standard Gaussian,

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}} e^{t(-\log \widetilde{W})} &= \mathbb{E}_{\mathbb{Q}} \widetilde{W}^{-t} = \mathbb{E}\left(W_0^{1-t} + W_1^{1-t}\right) \\ &= 2\,\mathbb{E}2^{-(1-t)}e^{(1-t)\sqrt{2\log 2}N - (1-t)\log 2} = 2^{2t-1+(1-t)^2} \\ &= e^{t^2 \log 2}. \end{aligned}$$

□

Define the measure ν on the positive real axis by setting

$$\int f \, \mathrm{d}\nu = \mathbb{E} Y f(Y) = \mathbb{E}_{\mathbb{Q}} \widetilde{Y} f(\widetilde{Y}) \tag{10}$$

for all continuous functions $f : \mathbb{R}^+ \to \mathbb{R}$ with compact support. The asymptotics of this measure will be determined through the functions

$$F_{\alpha,\beta}(x) = \nu((\alpha e^x, \beta e^x]) \quad \text{for } 0 < \alpha < \beta.$$

In terms of $F_{\alpha,\beta}$, the statement of Theorem 1 is essentially equivalent to the following proposition.

Proposition 6. *Let $F_{\alpha,\beta}$ be defined by ν as above. Then*

$$F_{\alpha,\beta}(x) \sim c_1 \log \frac{\beta}{\alpha} \quad \text{as } x \to \infty,$$

where

$$c_1 = \frac{2}{\log 2} \mathbb{E} \mu([0,1/2]) \log \left(1 + \frac{\mu([1/2,1])}{\mu([0,1/2])} \right) < \infty.$$

The first step towards the proof of the proposition above is deriving the Poisson equation satisfied by $F_{\alpha,\beta}$. Let τ denote the law of $-\log \widetilde{W}$. By using the independence of $\widetilde{W}$ and $\widetilde{Y}$ we get

$$\begin{aligned} \tau * F_{\alpha,\beta}(x) &= \int_{\mathbb{R}} \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\widetilde{Y} \in (\alpha e^{x+y}, \beta e^{x+y}]\}} \tau(\mathrm{d}y) \\ &= \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\widetilde{W}\widetilde{Y} \in (\alpha e^x, \beta e^x]\}} \\ &= F_{\alpha,\beta}(x) + \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\widetilde{W}\widetilde{Y} \in (\alpha e^x, \beta e^x]\}} - \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\widetilde{Y} \in (\alpha e^x, \beta e^x]\}}, \end{aligned}$$

where the convolution of the measure τ with a function $F : \mathbb{R} \to \mathbb{R}$ is defined by

$$\tau * F(x) = \int_{\mathbb{R}} F(x+y) \tau(\mathrm{d}y) = \int_{\mathbb{R}} F(x-y) \tau(\mathrm{d}y).$$

By using part (2) of Lemma 5, the distributional equation (9) and the definitions of the variables $\widetilde{W}$, $\widetilde{Y}$ and $\widetilde{B}$, the term $\mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\widetilde{Y} \in (\alpha e^x, \beta e^x]\}}$ above may be expressed as

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\widetilde{Y} \in (\alpha e^x, \beta e^x]\}} &= \mathbb{E} Y \, \mathbf{1}_{\{Y \in (\alpha e^x, \beta e^x]\}} \\ &= \mathbb{E} \Big(W_0 Y_0 \mathbf{1}_{\{\alpha e^x - W_1 Y_1 < W_0 Y_0 \leq \beta e^x - W_1 Y_1\}} \\ &\qquad + W_1 Y_1 \mathbf{1}_{\{\alpha e^x - W_0 Y_0 < W_1 Y_1 \leq \beta e^x - W_0 Y_0\}} \Big) \\ &= \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\alpha e^x - \widetilde{B} < \widetilde{W}\widetilde{Y} \leq \beta e^x - \widetilde{B}\}}. \end{aligned}$$

The previous computations imply that $F_{\alpha,\beta}$ satisfies the Poisson equation

$$F_{\alpha,\beta}(x) - \tau * F_{\alpha,\beta}(x) = \psi_{\alpha,\beta}(x) \tag{11}$$

with the function $\psi_{\alpha,\beta}$ given by

$$\psi_{\alpha,\beta}(x) = \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\alpha e^x - \widetilde{B} < \widetilde{W}\widetilde{Y} \leq \beta e^x - \widetilde{B}\}} - \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \, \mathbf{1}_{\{\alpha e^x < \widetilde{W}\widetilde{Y} \leq \beta e^x\}}. \tag{12}$$

The desired results on the solutions of this Poisson equation at infinity could be achieved almost exactly in the same way as in Buraczewski [12], that is, by building on the general theory developed by Port and Stone [36]. The following proposition is originally due to Buraczewski, but we prefer to give it a self-contained proof of independent interest that uses only elementary Fourier analysis.

Proposition 7. *Let ν be a locally finite (non-negative) Borel measure on $[0,\infty)$ that grows at most polynomially in the sense that there exist $\gamma, C > 0$ such that*

$$\nu((0,x]) \le C(1+x)^\gamma \quad \text{for all } x \ge 0.$$

Define the functions

$$F_{\alpha,\beta}(x) = \nu((\alpha e^x, \beta e^x]) \quad \text{for all } 0 < \alpha < \beta$$

and assume that for each α, β the function $\psi_{\alpha,\beta} : \mathbb{R} \to \mathbb{R}$ is a bounded and continuous function indexed by the parameters α and β that satisfies

$$\int_{-\infty}^{\infty} (1+|x|)^2 \, |\psi_{\alpha,\beta}(x)| \, \mathrm{d}x < \infty \quad \text{and}$$

$$\int_{-\infty}^{\infty} \psi_{\alpha,\beta}(x) \, \mathrm{d}x = 0.$$

Denote

$$C_{\alpha,\beta} = \int_{-\infty}^{\infty} x\psi_{\alpha,\beta}(x) \, \mathrm{d}x$$

and assume that the map $(\alpha,\beta) \mapsto C_{\alpha,\beta}$ is continuous. Finally, let τ be a Gaussian measure on $\mathbb{R}$, i.e. τ is the law of a centered Gaussian random variable with variance $\sigma^2 > 0$.

Then, if $F_{\alpha,\beta}$ satisfies the Poisson equation

$$F_{\alpha,\beta} - \tau * F_{\alpha,\beta} = \psi_{\alpha,\beta}$$

it has the asymptotics

$$\lim_{x\to\infty} F_{\alpha,\beta}(x) = \frac{2}{\sigma^2} C_{\alpha,\beta}.$$

We split our proof of this proposition into two lemmas and a finalizing convolution argument.

Lemma 8. *Let the function $F : \mathbb{R} \to \mathbb{R}$ be bounded from below and satisfy*

$$\lim_{x\to-\infty} F(x) = 0.$$

Assume also that F grows at most exponentially at infinity[1] *and solves the Poisson equation*

$$F - \tau * F = \psi, \tag{13}$$

where $\tau \sim N(0,\sigma^2)$ is as in Proposition 7 and $\psi : \mathbb{R} \to \mathbb{R}$ satisfies

$$\int_{-\infty}^{\infty} (1+|x|)^2 |\psi(x)| \, \mathrm{d}x < \infty, \quad \int_{-\infty}^{\infty} \psi(x) \, \mathrm{d}x = 0 \quad \text{and} \quad \lim_{x\to\pm\infty} \psi(x) = 0.$$

---

[1] The assumption of exponential growth is used only to ensure that the convolution with τ is well defined.

Then F has the asymptotics

$$\lim_{x\to\infty} F(x) = \frac{2}{\sigma^2}\int_{-\infty}^{\infty} x\psi(x)\,\mathrm{d}x. \tag{14}$$

Proof. To shorten the notations we denote $\int_{-\infty}^{\infty} x\psi(x)\,\mathrm{d}x = A$.

We start by proving that the equation (13) has some bounded solution F_1 that has the desired asymptotics

$$\lim_{x\to-\infty} F_1(x) = 0 \quad \text{and} \quad \lim_{x\to\infty} F_1(x) = \frac{2}{\sigma^2}A. \tag{15}$$

We first consider the case $A = 0$. Then our assumptions imply that the Fourier transform of ψ satisfies

$$\widehat{\psi} \in C^2(\mathbb{R}) \cap L^\infty(\mathbb{R}) \quad \text{and} \quad \widehat{\psi}(0) = \widehat{\psi}'(0) = 0.$$

As $1 - \widehat{\tau}(\xi) = 1 - \exp(-\sigma^2\xi^2/2)$ is smooth with zero of order 2 at the origin we may directly define F_1 through

$$\widehat{F_1}(\xi) := \frac{\widehat{\psi}(\xi)}{1 - e^{-\sigma^2\xi^2/2}} = \widehat{\psi}(\xi) + \left(\frac{\widehat{\psi}(\xi)}{\xi^2}\right)\left(\frac{\xi^2 e^{-\sigma^2\xi^2/2}}{1 - e^{-\sigma^2\xi^2/2}}\right) =: \widehat{\psi}(\xi) + \widehat{F_2}(\xi).$$

Since obviously $\widehat{F_2} \in L^1$, we have $\lim_{x\to\pm\infty} F_2(x) = 0$ by the Riemann-Lebesgue lemma, and the same follows for F_1 by the assumption on ψ.

In order to construct a solution F_1 in the general case $A \neq 0$, first define $\psi_0 = F_0 - \tau * F_0$ with $F_0 = \chi_{(0,\infty)}$. Directly from the definition we see that $\psi_0 \in L^\infty(\mathbb{R})$ and that ψ_0 decays exponentially as $x \to \pm\infty$, so that it satisfies the moment conditions of the lemma and moreover $\widehat{\psi_0} \in C^\infty$. Also, $\widehat{F_0}(\xi) = \pi\delta_0 - i\xi^{-1}$ (here ξ^{-1} is understood as a principal value distribution). Since $1 - \widehat{\tau}(\xi) = \sigma^2\xi^2/2 + O\left(\xi^3\right)$, we see that

$$\widehat{\psi_0}(\xi) = -\frac{i\sigma^2}{2}\xi + O(\xi^2)$$

at the origin. Hence

$$\int_{-\infty}^{\infty} \psi_0(x)\,\mathrm{d}x = \widehat{\psi_0}(0) = 0 \quad \text{and} \quad \int_{-\infty}^{\infty} x\psi_0(x)\,\mathrm{d}x = i\widehat{\psi}_0'(0) = \frac{\sigma^2}{2}.$$

Thus, in the case $A \neq 0$ we define F_1 by finding the solution for the Poisson equation (13) with the right hand side $\widetilde{\psi} = \psi - \frac{2}{\sigma^2}A\psi_0$ and then adding $\frac{2}{\sigma^2}AF_0$. At this point it is clear that the solution obtained this way is bounded and has the desired behaviour at $\pm\infty$.

Let us finally assume that F and ψ are as in the theorem. Let F_1 be the bounded solution of (13) constructed above, so that F_1 satisfies the conclusion of the theorem. It is enough to verify that $H := F - F_1$ is constant since then $H \equiv 0$ by considering the limit at $-\infty$. Now H is bounded from below and satisfies the homogeneous Poisson equation

$$H = \tau * H. \tag{16}$$

The claim follows from Lemma 9 below. □

Lemma 9. *Let H solve the homogeneous Poisson equation* (16) *and assume that it is bounded from below and has at most exponential growth at $\pm\infty$. Then H is constant.*

Proof. By adding a constant we may without loss of generality assume that $H \geq 0$. Let $u(x,t)$ ($x \in \mathbb{R}$, $t \geq 0$) denote the heat extension of H to the upper half-plane, explicitly given by

$$u(x,t) = \frac{1}{\sqrt{2\pi t}} \int_{-\infty}^{\infty} e^{\frac{-(y-x)^2}{2t}} H(y)\,\mathrm{d}y.$$

By assumption, u is periodic in t: denoting $t_0 = \sigma^2$, $u(x,t+t_0) = u(x,t)$ for all $t \geq 0$. Define the function v in the upper-half plane by setting

$$v(x,t) := \int_0^{t_0} u(x,t+s)\,\mathrm{d}s.$$

Then v solves the heat equation and, by the periodicity of u, it is constant in t. This it is harmonic in x, i.e. a linear function $v(x,t) = ax + b$. Here $a = 0$ by the nonnegativity of u, whence v is constant. This shows that there is a constant C independent of x so that $\int_{t_0/2}^{t_0} u(x,s)ds < C$. Especially, for each x there is $t_1 = t_1(x) \in (t_0/2, t_0)$ so that $u(x,t_1) \leq 2C/t_0$. The heat kernel $(2\pi t)^{-1/2}e^{-x^2/2t}$ can be bounded from below on $x \in [-1,1]$ uniformly in $t \in (t_0/2, t_0)$, whence again using the nonnegativity of H we deduce that $\int_{x-1}^{x+1} H(y)\,\mathrm{d}y \leq C'$ for all $x \in \mathbb{R}$. As we combine this information with the fact that $H = \tau * H$ it follows that H is bounded. Then the equation

$$(1 - e^{-\xi^2/2})\widehat{H}(\xi) = 0,$$

interpreted in the sense of distributions, shows that $\widehat{H} = c_1\delta_0 + c_2\delta_0'$, i.e. H is linear. By nonnegativity we finally deduce that H is constant. ☐

We finish the proof of Proposition 7 by deducing it from Lemma 8 by a convolution argument analogous to the one of Buraczewski [12].

Proof of Proposition 7. Let $\phi \geq 0$ be an arbitrary symmetric smooth test function with $\operatorname{supp}\phi \subset [-1,1]$ and $\int_R \phi = 1$. Given any locally integrable $g : \mathbb{R} \to \mathbb{R}$ let g_ε denote the convolution $g_\varepsilon = g * \varepsilon^{-1}\phi(\varepsilon^{-1}\cdot)$. By convolving the Poisson equation we obtain (writing e.g. $(F_{\alpha,\beta})_\varepsilon = F_{\alpha,\beta,\varepsilon}$) for any $0 \leq \alpha < \beta$ and $\varepsilon > 0$

$$F_{\alpha,\beta,\varepsilon} = \tau * F_{\alpha,\beta,\varepsilon} + \psi_{\alpha,\beta,\varepsilon}.$$

By the continuity of $\psi_{\alpha,\beta,\varepsilon}$ and integrability of $\psi_{\alpha,\beta}$ we have $\lim_{x\to\pm\infty} \psi_{\alpha,\beta,\varepsilon} = 0$. From Lemma 8 we thus obtain, for each $\varepsilon > 0$, the asymptotics

$$\lim_{x\to\pm\infty} F_{\alpha,\beta,\varepsilon}(x) = \frac{2}{\sigma^2}C_{\alpha,\beta}, \tag{17}$$

since

$$\int_{\mathbb{R}} x\psi_{\alpha,\beta,\varepsilon}(x)\,\mathrm{d}x = i\widehat{\psi}'_{\alpha,\beta,\varepsilon}(0) = i\widehat{\psi}'_{\alpha,\beta}(0) = \int_{\mathbb{R}} x\psi_{\alpha,\beta}(x) = C_{\alpha,\beta}.$$

In order to remove the ε from (17), let $k \in (1, (\beta/\alpha)^{1/2})$ be given and observe that by the definition of $F_{\alpha,\beta}$ as a measure of an interval we have $F_{k\alpha,k^{-1}\beta}(x) \leq F_{\alpha,\beta,\varepsilon}(x)$ for all x as soon as ε is small enough. Hence, we obtain from (17) that

$$\limsup_{x\to\infty} F_{\alpha,\beta}(x) \leq \frac{2}{\sigma^2}C_{k^{-1}\alpha,k\beta}.$$

By letting $k \to 1^+$ and recalling the assumption of the continuity of $(\alpha,\beta) \mapsto C_{\alpha,\beta}$ it follows that $\limsup_{x\to\infty} F_{\alpha,\beta}(x) \le 2C_{\alpha,\beta}/\sigma^2$. The converse direction $\liminf_{x\to\infty} F_{\alpha,\beta}(x) \ge 2C_{\alpha,\beta}/\sigma$ is obtained analogously by starting from the inequality $F_{k^{-1}\alpha,k\beta}(x) \le F_{\alpha,\beta,\varepsilon}(x)$. □

The proof of Proposition 6 has now essentially been reduced to checking that the Poisson equation (11) with $F_{\alpha,\beta}$ determined by ν as in (10) and $\psi_{\alpha,\beta}$ given by (12) satisfies the assumptions of Proposition 7.

Proof of Proposition 6. We first check that the measure ν satisfies $\nu((0,x]) \le C(1+x)^\gamma$ for some $C,\gamma > 0$. This is clear from the definition and Theorem A: for any $\gamma \in (0,1)$,

$$\nu((0,x]) = \mathbb{E}Y\mathbf{1}_{\{Y\in(0,x]\}} \le x^\gamma \mathbb{E}Y^{1-\gamma}\mathbf{1}_{\{Y\in(0,x]\}} \le \mathbb{E}Y^{1-\gamma}\,x^\gamma.$$

To check the integrability conditions on $\psi_{\alpha,\beta}$ we define the functions

$$\psi_\alpha(x) = \mathbb{E}_{\mathbb{Q}}\widetilde{Y}\,\mathbf{1}_{\left\{\alpha e^x - \widetilde{B} < \widetilde{W}\widetilde{Y} \le \alpha e^x\right\}} \quad \text{and} \quad \psi_\beta(x) = \mathbb{E}_{\mathbb{Q}}\widetilde{Y}\,\mathbf{1}_{\left\{\beta e^x - \widetilde{B} < \widetilde{W}\widetilde{Y} \le \beta e^x\right\}}.$$

By this definition,

$$\psi_{\alpha,\beta}(x) = \mathbb{E}_{\mathbb{Q}}\widetilde{Y}\,\mathbf{1}_{\{\alpha e^x - \widetilde{B} < \widetilde{W}\widetilde{Y} \le \beta e^x - \widetilde{B}\}} - \mathbb{E}_{\mathbb{Q}}\widetilde{Y}\,\mathbf{1}_{\widetilde{W}\widetilde{Y}\in(\alpha e^x,\beta e^x]} = \psi_\alpha(x) - \psi_\beta(x).$$

Since the functions ψ_α and ψ_β are positive, to check the integrability conditions of Theorem 7 on the functions $\psi_{\alpha,\beta}$ it is sufficient to show that

$$\int_{-1}^{1} \psi_\alpha(x)\,\mathrm{d}x < \infty \quad \text{and} \quad \int_{-\infty}^{\infty} x^2\psi_\alpha(x)\,\mathrm{d}x < \infty \quad \text{for all } \alpha > 0.$$

In our situation the first condition is trivial. For the second condition, some computation and a separate lemma are needed. We write

$$\mathbf{1}_{\left\{\alpha e^x - \widetilde{B} < \widetilde{W}\widetilde{Y} \le \alpha e^x\right\}} = \mathbf{1}_{\left\{\frac{\widetilde{W}\widetilde{Y}}{\alpha} \le t < \frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right\}} \quad \text{for } t = e^x$$

and use the integral

$$\int_a^b \frac{\log t}{t}\,\mathrm{d}t = \frac{1}{2}\Big(\log\frac{b}{a}\Big)\Big(\log ab\Big) \quad \text{for } 0 < a < b$$

to compute

$$
\begin{aligned}
&\int_0^\infty x^2 \psi_\alpha(x)\,\mathrm{d}x \\
&\quad = \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \int_0^\infty x^2 \mathbf{1}_{\{\alpha e^x - \widetilde{B} < \widetilde{W}\widetilde{Y} \le \alpha e^x\}}\,\mathrm{d}x \\
&\quad = \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \int_1^\infty \mathbf{1}_{\left\{\frac{\widetilde{W}\widetilde{Y}}{\alpha} \le t < \frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right\}} \frac{(\log t)^2}{t}\,\mathrm{d}t \\
&\quad = \mathbb{E}_{\mathbb{Q}} \widetilde{Y}\, \mathbf{1}_{\left\{\frac{\widetilde{W}\widetilde{Y}}{\alpha} > 1\right\}} \int_{\frac{\widetilde{W}\widetilde{Y}}{\alpha}}^{\frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}} \frac{(\log t)^2}{t}\,\mathrm{d}t \\
&\qquad + \mathbb{E}_{\mathbb{Q}} \widetilde{Y}\, \mathbf{1}_{\left\{\frac{\widetilde{W}\widetilde{Y}}{\alpha} < 1 < \frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right\}} \int_1^{\frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}} \frac{(\log t)^2}{t}\,\mathrm{d}t \\
&\quad \le \frac{1}{2} \mathbb{E}_{\mathbb{Q}} \widetilde{Y}\, \mathbf{1}_{\left\{\frac{\widetilde{W}\widetilde{Y}}{\alpha} > 1\right\}} \log\left(1 + \frac{\widetilde{B}}{\widetilde{W}\widetilde{Y}}\right) \log\left(\frac{\widetilde{W}\widetilde{Y}}{\alpha} \cdot \frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right) \log\left(\frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right) \\
&\qquad + \frac{1}{2} \mathbb{E}_{\mathbb{Q}} \widetilde{Y}\, \mathbf{1}_{\left\{\frac{\widetilde{W}\widetilde{Y}}{\alpha} < 1 < \frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right\}} \left(\log\left(\frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right)\right)^2 \log\left(\frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right) \\
&\quad =: I_1 + I_2,
\end{aligned}
$$

and similarly by the change of variables $s = e^{-x}$ we get

$$
\begin{aligned}
&\int_{-\infty}^0 x^2 \psi_\alpha(x)\,\mathrm{d}x \\
&\quad = \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \int_{-\infty}^0 x^2 \mathbf{1}_{\{\alpha e^x - \widetilde{B} < \widetilde{W}\widetilde{Y} \le \alpha e^x\}}\,\mathrm{d}x \\
&\quad = \mathbb{E}_{\mathbb{Q}} \widetilde{Y} \int_1^\infty \mathbf{1}_{\{\alpha/s - \widetilde{B} < \widetilde{W}\widetilde{Y} \le \alpha/s\}} \frac{(\log s)^2}{s}\,\mathrm{d}s \\
&\quad \le \frac{1}{2} \mathbb{E}_{\mathbb{Q}} \widetilde{Y}\, \mathbf{1}_{\left\{\frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha} < 1\right\}} \log\left(1 + \frac{\widetilde{B}}{\widetilde{W}\widetilde{Y}}\right) \log\left(\frac{\alpha}{\widetilde{W}\widetilde{Y}} \cdot \frac{\alpha}{\widetilde{W}\widetilde{Y}+\widetilde{B}}\right) \log\left(\frac{\alpha}{\widetilde{W}\widetilde{Y}}\right) \\
&\qquad + \frac{1}{2} \mathbb{E}_{\mathbb{Q}} \widetilde{Y}\, \mathbf{1}_{\left\{\frac{\widetilde{W}\widetilde{Y}}{\alpha} < 1 < \frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha}\right\}} \left(\log\left(\frac{\widetilde{W}\widetilde{Y}}{\alpha}\right)\right)^2 \log\left(\frac{\alpha}{\widetilde{W}\widetilde{Y}}\right) \\
&\quad =: I_3 + I_4.
\end{aligned}
$$

To show that $I_1 < \infty$ we use the crude estimate $\log(1+x) \le C_p x^p$, valid for all $p > 0$ for sufficiently large constant $C_p > 0$ depending only on p, to get

$$
\begin{aligned}
I_1 \le &\frac{C_{p_1} C_{p_2} C_{p_3}}{\alpha^{p_1} \alpha^{p_2}} \cdot \\
&\mathbb{E}\left(\mu([0,1/2]) \left(\frac{\mu([1/2,1])}{\mu([0,1/2])}\right)^{p_1} \left(\mu([0,1/2])^{p_2} + \mu([0,1])^{p_2}\right) \mu([0,1])^{p_3}\right).
\end{aligned}
$$

In Lemma 11 below we show that for any $0 < h < 1$ we have

$$
\mathbb{E}\mu([0,1/2])^h \mu([1/2,1])^h < \infty.
$$

By choosing $p_1, p_2, p_3 > 0$ such that $0 < 1 - p_1 + p_2 + p_3 < 1$ and $p_1 + p_2 + p_3 < 1$, this implies the finiteness of I_1. For I_2 one may estimate

$\frac{\widetilde{W}\widetilde{Y}+\widetilde{B}}{\alpha} \leq 1+\frac{\widetilde{B}}{\widetilde{W}\widetilde{Y}}$ and proceed as in the case of I_1. In estimating I_3 one may write $\frac{\alpha}{\widetilde{W}\widetilde{Y}+\widetilde{B}} < \frac{\alpha}{\widetilde{W}\widetilde{Y}}$ and proceed as before, and the finiteness of I_4 follows the same route.

In order to apply Proposition 7 we still need to show that

$$\int_{-\infty}^{\infty} \psi_{\alpha,\beta}(x)\,\mathrm{d}x = 0$$

and compute the value of the integral

$$C_{\alpha,\beta} = \int_{-\infty}^{\infty} x\psi_{\alpha,\beta}(x)\,\mathrm{d}x.$$

The first integral follows immediately from the integrability of ψ_α and the fact that

$$\psi_{\alpha,\beta}(x) = \psi_\alpha(x) - \psi_\beta(x) = \psi_\alpha(x) - \psi_\alpha(x + \log\frac{\alpha}{\beta}).$$

The value of $C_{\alpha,\beta}$ can be calculated by using the change of variables $x = e^t$ as above to obtain

$$\int_{-\infty}^{\infty} x\psi_\alpha(x)\,\mathrm{d}x = \frac{1}{2}\mathbb{E}_{\mathbb{Q}}\widetilde{Y}\log\left(1+\frac{\widetilde{B}}{\widetilde{W}\widetilde{Y}}\right)\log\frac{\widetilde{W}\widetilde{Y}(\widetilde{W}\widetilde{Y}+\widetilde{B})}{\alpha^2},$$

which implies

$$C_{\alpha,\beta} = \int_{-\infty}^{\infty} x(\psi_\alpha(x) - \psi_\beta(x))\,\mathrm{d}x = \log\frac{\beta}{\alpha}\,\mathbb{E}_{\mathbb{Q}}\widetilde{Y}\log\left(1+\frac{\widetilde{B}}{\widetilde{W}\widetilde{Y}}\right).$$

Proposition 7 now gives the desired asymptotics

$$\begin{aligned} F_{\alpha,\beta}(x) \quad &\overset{x\to\infty}{\longrightarrow} \quad \frac{2\,\mathbb{E}_{\mathbb{Q}}\widetilde{Y}\log\left(1+\frac{\widetilde{B}}{\widetilde{W}\widetilde{Y}}\right)}{2\log 2}\log\frac{\beta}{\alpha} \\ &= \quad \frac{2}{\log 2}\mathbb{E}\mu([0,1/2])\log\left(1+\frac{\mu([1/2,1])}{\mu([0,1/2])}\right)\log\frac{\beta}{\alpha}. \end{aligned}$$

for all $0 < \alpha < \beta$. □

Before moving on to the final step of the proof of Theorem 1 we complete the proof of Proposition 6 by proving Lemma 11.

Lemma 10. *For any $h \in (0,1)$ and any pair of intervals $I_1, I_2 \subset [0,1]$ such that $d(I_1, I_2) > 0$,*

$$\mathbb{E}\left(\mu(I_1)\mu(I_2)\right)^h < \infty.$$

Proof. Write

$$X_{t'}(x) = X_t(x) + (X_{t'}(x) - X_t(x))$$

for t large enough that so that for $t' > t$ one has

$$(X_{t'}(x) - X_t(x))_{x\in I_1} \perp (X_{t'}(x) - X_t(x))_{x\in I_2}.$$

Let $Z_1 = \sup_{x\in I_1\cup I_2} e^{\sqrt{2}X_t(x) - \mathbb{E}X_t(x)^2}$, $Z_2 = \sup_{x\in I_1\cup I_2} e^{-\sqrt{2}X_t(x) + \mathbb{E}X_t(x)^2}$, and

$$\mu^t(I_i) = \lim_{t'\to\infty}\sqrt{t}\int_{I_i} e^{\sqrt{2}(X_{t'}(x) - X_t(x)) - (\mathbb{E}X_{t'}(x)^2 - \mathbb{E}X_t(x)^2)}\,\mathrm{d}x.$$

Writing successively that $\mu(I_i) \le Z_1\mu^t(I_i)$ and $\mu^t(I_i) \le Z_2\mu(I_i)$ for $i = 1, 2$ we get

$$\begin{aligned}\mathbb{E}\left(\mu(I_1)\mu(I_2)\right)^h &\le (\mathbb{E}Z_1^{2h})(\mathbb{E}\mu^t(I_1)^h)(\mathbb{E}\mu^t(I_2)^h)\\ &\le (\mathbb{E}Z_1^{2h})(\mathbb{E}Z_2^h)^2(\mathbb{E}\mu(I_1)^h)(\mathbb{E}\mu(I_2)^h),\end{aligned}$$

and then Theorem A and standard estimates of Z_1 and Z_2 give the conclusion. □

Lemma 11. *For any* $h \in (0, 1)$,

$$\mathbb{E}\left(\mu([0,1/2])\mu([1/2,1])\right)^h < \infty.$$

Proof. Fix $h \in (0,1)$. For every $k \in \mathbb{N}$, let $J_k = \left[1/2 - 2^{-k}, 1/2 + 2^{-k}\right]$. Denote the left and right half of J_k by J_k^0 and J_k^1, and the right and left halves of J_k^0 (and J_k^1) by J_k^{00} and J_k^{01} (J_k^{10} and J_k^{11}). Define the sets A_k by

$$A_k = (J_k^{00} \times J_k^{11}) \cup (J_k^{00} \times J_k^{10}) \cup (J_k^{01} \times J_k^{11}).$$

Write

$$Z = \mu([0,1/2])\mu([1/2,1]) = \int_{[0,1/2]} \mu(\,\mathrm{d}x) \int_{[1/2,1]} \mu(\,\mathrm{d}y)$$

and define the random variables

$$\begin{aligned} Z_k &= \int_{[0,1/2]} \mu(\,\mathrm{d}x) \int_{[1/2,1]} \mu(\,\mathrm{d}y)\, \chi_{A_k}(x,y) \\ &= \mu(J_k^{00})\mu(J_k^{11}) + \mu(J_k^{00})\mu(J_k^{10}) + \mu(J_k^{01})\mu(J_k^{00}) \end{aligned}$$

for $k \in \mathbb{N}$. It is clear that

$$Z = \sum_{k=1}^{\infty} Z_k$$

and thus by the subadditivity of $x \mapsto x^h$

$$\mathbb{E}Z^h \le \sum_{k=1}^{\infty} \mathbb{E}Z_k^h.$$

By the exact scaling property of the construction the measure μ satisfies

$$\left(\mu(J_k^{\sigma_1\sigma_2})\right)_{\sigma_1,\sigma_2\in\{0,1\}} = 2^{-k+1}e^{\sqrt{2}X(J_k)-\mathbb{E}X(J_k)^2}\left(\mu'(J_1^{\sigma_1\sigma_2})\right)_{\sigma_1,\sigma_2\in\{0,1\}}, \tag{18}$$

where $X(J_k) = W(\mathcal{C}(J_k))$ is a centered Gaussian random variable with variance $\lambda(\mathcal{C}(J_k)) = (k-1)\log 2$ and μ' is random measure independent of $X(J_k)$ that has the same distribution as μ. But this implies that

$$Z_k \overset{d}{=} 2^{-2k+2}e^{2\sqrt{2}X(J_k)-2\mathbb{E}X(J_k)^2}Z_1',$$

where $Z_1' \overset{d}{=} Z_1$ is a random variable independent of Z_1. Since

$$2^{(-2k+2)h}\mathbb{E}e^{2\sqrt{2}hX(J_k)-2h\mathbb{E}X(J_k)^2} = 2^{(-2k+2)h}2^{(4h^2-2h)(k-1)} = 2^{4(h^2-h)(k-1)}$$

and $\mathbb{E}Z_1^h$ is finite by Lemma 10, we have

$$\mathbb{E}Z^h \le \mathbb{E}Z_1^h \sum_{k=1}^{\infty} 2^{4(h^2-h)(k-1)} < \infty.$$

□

Proof of Theorem 1. We will show that for any $r > 1$ there exists a λ_r such that

$$\mathbb{P}(Y > \lambda) \leq \frac{c_1}{\lambda} r \quad \text{for all } \lambda \geq \lambda_r.$$

The verification of the lower bound is similar and is left to the reader.

Let $r > 1$ and fix $q > 1$ so that $\frac{q \log q}{q-1} < \sqrt{r}$. By Proposition 6 there exists a λ_r such that

$$F_{1,q}(x) \leq c_1 \sqrt{r} \log q \quad \text{for all } x \geq \log \lambda_r.$$

By the definition $F_{1,q}(x) = \mathbb{E}(Y \mathbf{1}_{\{Y \in e^x, qe^x\}})$ we now have for $\lambda \geq \lambda_r$

$$\begin{aligned} \mathbb{P}(Y > \lambda) &= \sum_{k=0}^{\infty} \mathbb{P}\left(Y \in \left(\lambda q^k, \lambda q^{k+1}\right]\right) \leq \frac{1}{\lambda} \sum_{k=0}^{\infty} q^{-k} F_{1,q}(k \log q + \log \lambda) \\ &\leq \frac{1}{\lambda} \sum_{k=0}^{\infty} q^{-k} c_1 \sqrt{r} \log q = \frac{c_1}{\lambda} \sqrt{r} \frac{q \log q}{q-1} \leq \frac{c_1}{\lambda} r, \end{aligned}$$

as was to be shown. □

3. MODULUS OF CONTINUITY

In this section we prove Theorem 2. Our plan of attack is to follow the arguments carried out in [10] in the case of multiplicative cascades. However, the delicate dependence structure of multiplicative chaos calls for non-trivial modifications.

Let $((X_t(x))_{x \in \mathbb{R}})_{t \geq 0}$ be the exactly scale invariant Gaussian field on $\mathbb{R}$ as before and define the Gaussian field $((Y_t(x))_{x \in \mathbb{R}})_{t \geq 0}$ by setting

$$Y_t(x) = W(\mathcal{C}_t(x) \setminus \mathcal{C}_0(x)) = X_t(x) - X_0(x) \quad \text{for } x \in \mathbb{R}, t \geq 0.$$

In the proof of Theorem 2 it is convenient to use the characterization (7) of critical lognormal multiplicative chaos. To keep the notation simpler, we normalize the construction by the deterministic constant $c > 0$ in (7). Explicitly, we consider the critical measures associated to the fields X and Y and denote

$$\begin{aligned} \mu_{\sqrt{2}}(\mathrm{d}x) &= \lim_{t \to \infty} \sqrt{t} e^{\sqrt{2} X_t(x) - (t+1)} \, \mathrm{d}x \quad \text{and} \\ \nu_{\sqrt{2}}(\mathrm{d}x) &= \lim_{t \to \infty} \sqrt{t} e^{\sqrt{2} Y_t(x) - t} \, \mathrm{d}x, \end{aligned}$$

where the limits exist in probability in the weak sense. By construction it is clear that almost surely, the Radon-Nikodym derivative $\frac{\mathrm{d}\mu_{\sqrt{2}}}{\mathrm{d}\nu_{\sqrt{2}}}(x) = e^{\sqrt{2} X_0(x) - 1}$ is almost surely positive and uniformly bounded away from 0 and ∞ for all $x \in [0, 1]$, so for the purpose of our result on the modulus of continuity the difference between these two measures is insignificant. The measure $\mu_{\sqrt{2}}$ is exactly scale invariant as before, but in this section we make more use of the measure $\nu_{\sqrt{2}}$ which satisfies the $\star$-*scaling relation*: for every $\epsilon \in (0, 1]$ we have

$$(\nu_{\sqrt{2}}(A))_{A \in \mathcal{B}(\mathbb{R})} \overset{d}{=} \left(\int_A e^{\sqrt{2} Y_{-\log \epsilon} + 2 \log \epsilon} \nu^{\epsilon}_{\sqrt{2}}(\mathrm{d}x) \right)_{A \in \mathcal{B}(\mathbb{R})}, \tag{19}$$

where $\nu^{\epsilon}_{\sqrt{2}}$ is independent of $Y_{-\log\epsilon}$ and $(\nu^{\epsilon}_{\sqrt{2}}(A))_A \overset{d}{=} (\nu_{\sqrt{2}}(\epsilon^{-1}A))_A$. The proof of this scaling relation is recalled in the Appendix.

In the proof of the theorem we need bounds for the tail probabilities of $\mu_{\sqrt{2}}(I)$ and $\nu_{\sqrt{2}}(I)$ in terms of the length of the interval I. For the measure $\mu_{\sqrt{2}}$ the bounds will follow from the exact scaling relation and Theorem 1, so we begin by translating the upper bound given by Theorem 1 for the measure $\nu_{\sqrt{2}}$.

Lemma 12. *There exists a constant $C > 0$ such that for $x > 0$*

$$\mathbb{P}(\nu_{\sqrt{2}}([0,1]) > x) \le \frac{C}{x}. \tag{20}$$

Proof. The covariance of X_0 is $\mathbb{E}(X_0(x)X_0(y)) = \max(0, 1-|x-y|)$ and for every $t \ge 0$, $(X_0(x))_{x\in[0,1]}$ and $(Y_t(x))_{x\in[0,1]}$ are independent. We thus have

$$\begin{aligned}
\mathbb{P}(\mu_{\sqrt{2}}([0,1]) > \lambda) &= \mathbb{P}\Big(\int_0^1 e^{\sqrt{2}X_0(x)-1}\nu_{\sqrt{2}}(\,\mathrm{d}x) > \lambda\Big) \\
&\ge \mathbb{P}(e^{\sqrt{2}\min_{x\in[0,1]} X_0(x)}\nu_{\sqrt{2}}([0,1]) > e\lambda) \\
&\ge \mathbb{P}(\nu_{\sqrt{2}}([0,1]) > e\lambda,\ \min_{x\in[0,1]} X_0(x) > 0) \\
&= \mathbb{P}(\nu_{\sqrt{2}}([0,1]) > e\lambda)\mathbb{P}(\min_{x\in[0,1]} X_0(x) > 0).
\end{aligned}$$

Since $\mathbb{P}(\min_{x\in[0,1]} X_0(x) > 0) > 0$, Theorem 1 implies the existence of a constant $C > 0$ such that $\mathbb{P}(\nu_{\sqrt{2}}([0,1]) > \lambda) \le C\lambda^{-1}$.

□

We shall also need the following auxiliary lemma.

Lemma 13. *Let $X \ge 0$ satisfy $\mathbb{P}(X > \lambda) \le \dfrac{A}{\lambda}$ for $\lambda > 0$.*

Let X_j, $j \in \{1,\dots,N\}$ be (possibly dependent) random variables with the same distribution as X and let $a_j \ge 0$ for $j \in \{1,...,N\}$. Then

$$\mathbb{P}\big(\sum_{j=1}^N a_jX_j > \lambda\big) \le \frac{C\cdot A\log(N+1)\big(\sum_{j=1}^N a_j\big)}{\lambda} \qquad \text{for all } \ \lambda > 0,$$

with a universal (in particular, independent of A) constant $C < \infty$.

Proof. We may assume that $\sum_{j=1}^N a_j = 1$ since the statement scales in the right way. Fix $t \in (0,1)$ and observe first that for all positive $y_1,\dots,y_N$ one has

$$\big(\sum_{j=1}^N a_jy_j\big)^t \le \sum_{j=1}^N a_j^ty_j^t$$

Fix $\lambda > 0$. The above holds if we set $y_j = (x_j-\lambda)_+$, where we denote the positive part by $y_+ := \max(0,y)$ and let, for now, the x_j:s be arbitrary real numbers. We obtain using $\sum_{j=1}^N a_j = 1$ that

$$\big(\sum_{j=1}^N a_jx_j - \lambda\big)_+^t \ \le\ \big(\sum_{j=1}^N a_j(x_j-\lambda)_+\big)^t \ \le\ \sum_{j=1}^N a_j^t(x_j-\lambda)_+^t,$$

or, in other words,

$$\phi(\sum_{j=1}^N a_j x_j) \le \sum_{j=1}^N a_j^t \phi(x_j),$$

where $\phi(x) := (x-\lambda)_+^t$. Especially, we have

$$\mathbb{E}\phi(\sum_{j=1}^N a_j X_j) \ \le \ \mathbb{E}\phi(X)\sum_{j=1}^N a_j^t. \tag{21}$$

The right hand side can be estimated as follows:

$$\begin{aligned}
\mathbb{E}\phi(X) \ &= \ \int_0^\infty \phi'(u)\mathbb{P}(X>u)\,du \ \le \ A\int_\lambda^\infty t(u-\lambda)^{t-1}u^{-1}\,du \\
&= \ At\int_0^\lambda \frac{y^{t-1}dy}{y+\lambda} + At\int_\lambda^\infty \frac{y^{t-1}dy}{y+\lambda} \\
&\le \ At\lambda^{-1}\int_0^\lambda y^{t-1}dy + At\int_\lambda^\infty y^{t-2}dy \\
&= \ A(1-t)^{-1}\lambda^{t-1}. \qquad (22)
\end{aligned}$$

We thus obtain from (21)

$$\big(\lambda\phi'(2\lambda)\big)\cdot\mathbb{P}(\sum_{j=1}^N a_j X_j > 2\lambda) \ \le \ \mathbb{E}\phi(\sum_{j=1}^N a_j X_j) \ \le \ \big(\sum_{j=1}^N a_j^t\big)\mathbb{E}\phi(X),$$

and by combining with (22)

$$\mathbb{P}(\sum_{j=1}^N a_j X_j > 2\lambda) \le \frac{A}{\lambda}\Big(\frac{1}{t(1-t)}\sum_{j=1}^N a_j^t\Big). \tag{23}$$

Finally, choosing $t = t_0 := 1 - 1/\log N$ (for $N \ge 3$) we get

$$\Big(\frac{1}{t_0(1-t_0)}\sum_{j=1}^N a_j^{t_0}\Big) \le \frac{N^{1-t_0}}{t_0(1-t_0)}\big(\sum_{j=1}^N a_j\big)^{t_0} = \frac{N^{1-t_0}}{t_0(1-t_0)} = et_0^{-1}\log N,$$

and then (23) yields the stated result. □

The next lemma contains the key technical estimates that lead to the proof of Theorem 2.

Lemma 14. *Let us index by $\sigma \in \Sigma_n = \{0,1\}^n$ the dyadic subintervals I_σ of $[0,1]$ of length 2^{-n}. Moreover, write $\Sigma_n^{(e)}$ for the family of even dyadic intervals of length 2^{-n} (i.e. intervals of the form $[(2j)2^{-n},(2j+1)2^{-n})$) and $\Sigma_n^{(o)}$ for the odd ones. Then for $\gamma \in (0,\frac{1}{2})$ and $\epsilon \in (0,1)$ there exists a constant $C = C(\epsilon)$ such that*

$$\mathbb{P}(\max_{\sigma\in\Sigma_n^{(e/o)}} \nu_{\sqrt{2}}(I_\sigma) \ge n^{-\gamma}) \le Cn^{(1-\epsilon)(\gamma-\frac{1}{2})} \tag{24}$$

Proof. We begin by noting that by specializing the $\star$-scaling relation to dyadics, we get

$$(\nu_{\sqrt{2}}(I_\sigma))_{\sigma\in\Sigma_n} \overset{d}{=} \left(2^{-n}\int_{I_\sigma} e^{\sqrt{2}Y_{n\log 2}(x)-n\log 2}\nu_{\sqrt{2}}^{(n)}(\,\mathrm{d}x)\right)_{\sigma\in\Sigma_n}, \tag{25}$$

where $\nu^{(n)}_{\sqrt{2}}$ is independent of $Y_{n\log 2}$ and $(\nu^{(n)}_{\sqrt{2}}(A))_A \overset{d}{=} (\nu_{\sqrt{2}}(2^n A))_A$. Since $Y_t(x)$ and $Y_t(y)$ are independent when $|x-y| \geq 1$, $\nu_{\sqrt{2}}(A)$ is independent of $\nu_{\sqrt{2}}(B)$ when $d(A,B)\geq 1$. Thus the scaling property implies that $(\nu^{(n)}_{\sqrt{2}}\lfloor I_\sigma)_{\sigma\in\Sigma_n^{(e)}}$ is a family of independent random measures (and similarly for the odd intervals) – here $\nu^{(n)}_{\sqrt{2}}\lfloor I_\sigma$ denotes the restriction of $\nu^{(n)}_{\sqrt{2}}$ to I_σ.

Let us write

$$W_{n,\sigma} = 2^{-n}\int_{I_\sigma} e^{\sqrt{2}Y_{n\log 2}(x)-n\log 2}\nu^{(n)}_{\sqrt{2}}(\,\mathrm{d}x). \tag{26}$$

Using the independence noted above, we see that

$$\begin{aligned}\mathbb{P}\left(\max_{\sigma\in\Sigma_n^{(e)}} W_{n,\sigma} < n^{-\gamma}\right) &= \mathbb{E}\prod_{\sigma\in\Sigma_n^{(e)}}\mathbb{P}(W_{n,\sigma}<n^{-\gamma}|Y_{n\log 2})\\ &\geq \mathbb{E}\prod_{\sigma\in\Sigma_n}(1-\mathbb{P}(W_{n,\sigma}\geq n^{-\gamma}|Y_{n\log 2}))\end{aligned} \tag{27}$$

To estimate $\mathbb{P}(W_{n,\sigma}\geq n^{-\gamma}|Y_{n\log 2})$, we will approximate the integral by a Riemann sum and then make use of the previous lemma. We will need to consider $\mathbb{P}(\nu^{(n)}_{\sqrt{2}}([0,2^{-n-k}])>\lambda)$, where the auxiliary index $k=k_n$ will be chosen later on. Using the definition of $\nu^{(n)}_{\sqrt{2}}$ and a similar argument as when estimating $\mathbb{P}(\nu_{\sqrt{2}}([0,1])>\lambda)$, we find

$$\mathbb{P}(\nu^{(n)}_{\sqrt{2}}([0,2^{-n-k}])>\lambda) \leq \frac{\mathbb{P}(\mu_{\sqrt{2}}([0,2^{-k}])>e\lambda)}{\mathbb{P}(\min_{x\in[0,2^{-k}]}X_0(x)>0)}.$$

Note that we can bound the denominator from below uniformly in k (for example by $\mathbb{P}(\min_{x\in[0,1]}X_0(x)>0)>0$). Using the exact scaling of $\mu_{\sqrt{2}}$, we see that

$$\mathbb{P}(\mu_{\sqrt{2}}([0,2^{-k}])>e\lambda) = \mathbb{P}(2^{-k}e^{\sqrt{2}X_k-\mathbb{E}(X_k^2)}\mu_{\sqrt{2}}([0,1])>e\lambda),$$

where X_k is a centered Gaussian independent of $\mu_{\sqrt{2}}$. Conditioning on X_k and using the tail of the distribution of $\mu_{\sqrt{2}}([0,1])$, we see that

$$\mathbb{P}(\mu_{\sqrt{2}}([0,2^{-k}])>e\lambda) \leq C2^{-k}\lambda^{-1}\mathbb{E}(e^{\sqrt{2}X_k-\mathbb{E}(X_k^2)}) = C2^{-k}\lambda^{-1}. \tag{28}$$

Collecting these arguments, we find that $\mathbb{P}(\nu^{(n)}_{\sqrt{2}}([0,2^{-n-k}])>\lambda)\leq C2^{-k}\lambda^{-1}$, where $C>0$ is independent of n,k and λ.

To get back to the size of $W_{n,\sigma}$, we need to consider a regularization of the function $f^{(n)}_{\sqrt{2}} := e^{\sqrt{2}Y_{n\log 2}-n\log 2}$ on the interval I_σ. Fix $\sigma\in\Sigma_n$ for the moment, let $k\in\mathbb{N}_+$ and divide I_σ into 2^k subintervals $(I_{\sigma,j})_{j=1}^{2^k}$ of equal length. Denote the midpoint of $I_{\sigma,j}$ by $x_{\sigma,j}$. Let $s>0$ and define the event $\mathcal{D}_s = \big\{\sup_{x\in I_{\sigma,j}}|Y_{n\log 2}(x)-Y_{n\log 2}(x_{\sigma,j})|\leq s \text{ for all } j=1,2,\ldots,2^k\big\}$. We then have on $\mathcal{D}_s$

$$\int_{I_\sigma} f^{(n)}_{\sqrt{2}}(x)\nu^{(n)}_{\sqrt{2}}(\,\mathrm{d}x) \leq e^{2\sqrt{2}s}\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x\,\nu^{(n)}_{\sqrt{2}}(I_{\sigma,j})$$

Let $\mathcal{F}_n = \sigma(\{Y_t(x) : x \in [0,1],\, t \leq n\log 2\})$. Since $\nu^{(n)}_{\sqrt{2}}(I_{\sigma,j}) \overset{d}{=} \nu^{(n)}_{\sqrt{2}}(I_{\sigma,i})$ for $j \neq i$ and the function $f^{(n)}_{\sqrt{2}}$ is independent of the measure $\nu^{(n)}_{\sqrt{2}}$, Lemma 13 and our estimate for $\mathbb{P}(\nu_{\sqrt{2}}(I_{\sigma,j}) > \lambda)$ imply that on $\mathcal{D}_s$

$$
\begin{aligned}
&\mathbb{P}\left(\int_{I_\sigma} f^{(n)}_{\sqrt{2}}(x)\nu^{(n)}_{\sqrt{2}}(\,\mathrm{d}x) > \lambda \,\middle|\, \mathcal{F}_n\right)\\
&\quad = \mathbb{P}\left(\frac{\int_{I_\sigma} f^{(n)}_{\sqrt{2}}(x)\nu^{(n)}_{\sqrt{2}}(\,\mathrm{d}x)}{e^{2\sqrt{2}s}\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x} > \frac{\lambda}{e^{2\sqrt{2}s}\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x} \,\middle|\, \mathcal{F}_n\right)\\
&\quad \leq \mathbb{P}\left(\frac{e^{2\sqrt{2}s}\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x\,\nu^{(n)}_{\sqrt{2}}(I_{\sigma,j})}{e^{2\sqrt{2}s}\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x} > \frac{\lambda}{e^{2\sqrt{2}s}\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x} \,\middle|\, \mathcal{F}_n\right)\\
&\quad \leq Ck2^{-k}\left(\frac{e^{2\sqrt{2}s}\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x}{\lambda}\right)
\end{aligned}
$$

for some constant $C > 0$. Setting $\lambda = n^{-\gamma}2^n$ and combining this inequality with (27) and the inequality $e^{-2x} \leq 1 - x$ valid for $x \in [0, 1/2]$, we get

$$
\begin{aligned}
&\mathbb{E}\mathbb{P}\left(\max_{\sigma\in\Sigma_n^{(e)}} \nu_{\sqrt{2}}(I_\sigma) > n^{-\gamma} \,\middle|\, \mathcal{F}_n\right)\\
&\qquad\qquad \leq 1 - \mathbb{E}\prod_{\sigma\in\Sigma_n}\left(1 - \mathbb{P}(W_{n,\sigma} \geq n^{-\gamma} | Y_{n\log 2})\right)\\
&\qquad\qquad \leq 1 - \mathbb{E}\exp\left(-2Ck2^{-k}e^{2\sqrt{2}s}\sum_{\sigma\in\Sigma_n}\left(\frac{\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x}{2^n n^{-\gamma}}\right)\right)\\
&\qquad\qquad\quad + 1 - \mathbb{P}(\mathcal{A}_{n,k,s})\\
&\qquad\qquad = 1 - \mathbb{E}\exp\left(-2Cke^{2\sqrt{2}s}n^{\gamma}\int_0^1 f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x\right) + 1 - \mathbb{P}(\mathcal{A}_{n,k,s})
\end{aligned}
$$

where $\mathcal{A}_{n,k,s}$ is the event

$$
\begin{aligned}
\mathcal{A}_{n,k,s} = &\left\{\max_{\sigma\in\Sigma_n} Ck2^{-k}\frac{e^{2\sqrt{2}s}\sum_{j=1}^{2^k}\fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x}{n^{-\gamma}2^n} < \frac{1}{2}\right\}\\
&\cap\left\{\sup_{x\in I_{\sigma,j}} |Y_{n\log 2}(x) - Y_{n\log 2}(x_{\sigma,j})| \leq s\,\forall j \in \{0,1,\ldots,2^k-1\}\,\forall\sigma\in\Sigma_n\right\}.
\end{aligned}
$$

Denoting

$$
S_n = n^{\frac{1}{2}}\int_0^1 e^{\sqrt{2}Y_{n\log 2}(x) - n\log 2}\,\mathrm{d}x,
$$

we finally get

$$
\begin{aligned}
\mathbb{P}\left(\max_{\sigma\in\Sigma_n^{(e)}} \mu_{\sqrt{2}}(I_\sigma) > n^{-\gamma}\right) &\leq 1 - \mathbb{E}\exp\left(-2Ce^{2\sqrt{2}s}kn^{(\gamma-\frac{1}{2})}S_n\right)\\
&\quad + 1 - \mathbb{P}(\mathcal{A}_{n,k,s}).
\end{aligned}
\tag{29}
$$

We then estimate the terms in the inequality above. Denote

$$\begin{aligned}\mathcal{B}_n &= \left\{ \max_{\sigma\in\Sigma_n} Ck2^{-k} \frac{e^{2\sqrt{2}s} \sum_{j=1}^{2^k} \fint_{I_{\sigma,j}} f^{(n)}_{\sqrt{2}}(x)\,\mathrm{d}x}{n^{-\gamma}2^n} < \frac{1}{2} \right\} \\ &= \{\max_{\sigma\in\Sigma_n} \int_{I_\sigma} e^{\sqrt{2}Y_{n\log 2}(x)-n\log 2}\,\mathrm{d}x < n^{-\gamma}(2Cke^{2\sqrt{2}s})^{-1}\}\end{aligned}$$

and

$$\mathcal{B}'_{n,k,s} = \left\{ \sup_{x\in I_{\sigma,j}} |Y_{n\log 2}(x) - Y_{n\log 2}(x_{\sigma,j})| \le s\,\forall j \in \{0,1,\dots,2^k-1\}\,\forall\sigma\in\Sigma_n \right\}$$

so that

$$\mathcal{A}_{n,k,s} = \mathcal{B}_n \cap \mathcal{B}'_{n,k,s} \quad\text{and}\quad 1-\mathbb{P}(\mathcal{A}_{n,k,s}) \le \big(1-\mathbb{P}(\mathcal{B}_n)\big) + \big(1-\mathbb{P}(\mathcal{B}'_{n,k,s})\big).$$

We first estimate the probability of $\mathcal{B}'_{n,k,s}$ not occurring. For all σ and j, the length of $I_{\sigma,j}$ is 2^{-n-k} and the covariance of the process $(Y_{n\log 2}(x))_{x\in I_{\sigma,j}}$ is 2^n-Lipschitz, so by the Borell–Tsirelson inequality we have, for any $\sigma\in\Sigma_n$ and $j=1,\dots,2^k$,

$$\mathbb{P}\left(\sup_{x\in I_{\sigma,j}} |Y_{n\log 2}(x) - Y_{n\log 2}(x_{\sigma,j})| > s\right) \le c(1+2^{\frac{k}{2}}s)e^{-2^{k-1}s^2},$$

where $c>0$ is an absolute constant. It follows that

$$1-\mathbb{P}(\mathcal{B}'_{n,k,s}) \le c\,2^{n+k}(1+2^{\frac{k}{2}}s)e^{-2^{k-1}s^2}.$$

For the choice $s_n \sim \sqrt{\epsilon\log n}$ and $k_n \sim \alpha\log n$, the right side of this estimate is asymptotically equivalent to

$$n^{\alpha\log 2}\left(1+n^{\frac{\alpha\log 2}{2}}\sqrt{\epsilon\log n}\right)e^{n\log 2-\frac{\epsilon}{2}n^{\alpha\log 2}\log n},$$

from which we see that in order to have $\sum_{n=1}^{\infty}(1-\mathbb{P}(\mathcal{B}'_{n,k_n,s_n}))<\infty$ we may take $\epsilon>0$ arbitrarily small, but must restrict to $\alpha\ge 1/\log 2$. Taking $\alpha = 1/\log 2$, in (29) these choices give

$$\begin{aligned}\mathbb{P}\left(\max_{\sigma\in\Sigma_n^{(e)}} \nu_{\sqrt{2}}(I_\sigma) > n^{-\gamma}\right) &\le 1-\mathbb{E}\exp\left(-2Ce^{2\sqrt{2}\sqrt{\epsilon\log n}}n^{(\gamma-\frac{1}{2})}S_n\right) \\ &\quad + c'n^{-c''\log n} + \big(1-\mathbb{P}(\mathcal{B}_n)\big)\end{aligned}$$

for some constants $c', c''>0$ depending on ϵ.

To estimate the probability of $\mathcal{B}_n$, we note that

$$\{S_n < n^{\frac{1}{2}-\gamma}(2Ck_ne^{2\sqrt{2}s_n})^{-1}\} \subset \mathcal{B}_n. \tag{30}$$

By Chebyshev's inequality we then see that for any $q<1$

$$\begin{aligned}1-\mathbb{P}(\mathcal{B}_n) &\le \mathbb{P}(S_n > (2Ck_ne^{2\sqrt{2}s_n})^{-1}n^{(\frac{1}{2}-\gamma)}) \\ &\le (2Ck_ne^{2\sqrt{2}s_n})^q\frac{\mathbb{E}(S_n^q)}{n^{(\frac{1}{2}-\gamma)q}}.\end{aligned}$$

If we knew that $\mathbb{E}(S_n^q)$ were uniformly bounded in n for some values of q, we would have a quantitative estimate for the speed at which $\mathbb{P}(\mathcal{B}_n)$ tends to one. For this we make use of Kahane's convexity inequalities [23]:

Proposition 15. *Let* $G : [0,\infty) \to \mathbb{R}$ *be a concave function such that* $|G(x)| \leq C(1+x^\alpha)$ *for some positive constants* C *and* α*. Let* $A \subset \mathbb{R}^d$ *be a Borel set,* ρ *be a Radon measure on* A *and* $(X_r)_{r\in A}$ *and* $(Y_r)_{r\in A}$ *be two continuous and centered Gaussian processes on* A *such that the covariance kernels satisfy* $k_X(u,v) \leq k_Y(u,v)$ *for all* $u, v \in A$*. Then*

$$\mathbb{E}\, G\left(\int_A e^{X_r - \frac{1}{2}\mathbb{E}(X_r^2)} \rho(\,\mathrm{d}r)\right) \geq \mathbb{E}\, G\left(\int_A e^{Y_r - \frac{1}{2}\mathbb{E}(Y_r^2)} \rho(\,\mathrm{d}r)\right).$$

To apply this inequality, we construct a Gaussian field on $[0,1]$ for which the moments of the corresponding measure can be calculated and for which we have a covariance structure that allows comparing with the field $Y_{n\log 2}$ (such a comparison is also used in [18] to prove that the limit of the total mass martingale associated to non-renormalized critical chaos measures vanishes almost surely).

The Gaussian field we shall employ is essentially a Gaussian branching random walk. Let us associate to the collection $\{I_\sigma\}$ of dyadic subintervals of $[0,1]$ an i.i.d. collection of standard Gaussian random variables $\{V_\sigma\}$. Define the field

$$U_n(x) = \sum_{k=1}^n \sum_{\sigma\in\Sigma_k:\ x\in I_\sigma} V_\sigma.$$

The covariance of U_n is given by

$$\begin{aligned}
\mathbb{E}(U_n(x)U_n(y)) &= \sum_{k,k'=1}^n \sum_{\sigma\in\Sigma_k \sigma'\in\Sigma_{k'}: x\in I_\sigma, y\in I_{\sigma'}} \mathbb{E}(V_\sigma V_{\sigma'}) \\
&= \sum_{k,k'=1}^n \sum_{\sigma\in\Sigma_k, \sigma'\in\Sigma_{k'}: x\in I_\sigma, y\in I_{\sigma'}} \mathbf{1}(\sigma=\sigma') \\
&= \sum_{k=1}^n \sum_{\sigma\in\Sigma_k:\ x,y\in I_\sigma} 1.
\end{aligned}$$

To compare this with the translation invariant field, we note that certainly to have a $\sigma \in \Sigma_k$ such that $x, y \in I_\sigma$, one must have $2^{-k} \geq |x-y|$. Thus

$$\begin{aligned}
\mathbb{E}(U_n(x)U_n(y)) &\leq \sum_{k=1}^{\frac{-\log|x-y|}{\log 2}\vee n} 1 \\
&= \frac{-\log|x-y|}{\log 2} \vee n \\
&\leq \frac{1}{\log 2}\mathbb{E}(Y_{n\log 2}(x)Y_{n\log 2}(y)) + C,
\end{aligned}$$

for some large enough constant C, since the covariance of the field $Y_{n\log 2}$ is given by

$$\mathbb{E}(Y_{n\log 2}(x)Y_{n\log 2}(y)) = \begin{cases} -\log|x-y| + |x-y| - 1, & 2^{-n} \leq |x-y| \leq 1 \\ n\log 2 + |x-y| - 2^n|x-y|, & |x-y| \leq 2^{-n} \end{cases}.$$

Let us thus consider a standard Gaussian variable Z independent of $Y_{n\log 2}$ and define the fields

$$A(x)=\sqrt{2\log 2}U_n(x) \quad \text{and} \quad B(x)=\sqrt{2}Y_{n\log 2}(x)+\sqrt{2C\log 2}Z.$$

We have $\mathbb{E}(A(x)A(y))\leq \mathbb{E}(B(x)B(y))$ for all x,y. We then apply the convexity inequality to the fields A and B with the convex function $G(x)=n^{q\frac{1}{2}}x^q$ for $q<1$, to get

$$\mathbb{E}(e^{q\sqrt{2C\log 2}Z-qC\log 2})\mathbb{E}(S_n^q)$$
$$\leq \mathbb{E}(n^{q\frac{1}{2}}(\int_0^1 e^{\sqrt{2\log 2}U_n(x)-\log 2\mathbb{E}(U_n(x)^2)}\,\mathrm{d}x)^q).$$

Comparing with the notation of [10], we see that the quantity on the right here is simply $\mathbb{E}((n^{\frac{1}{2}}Z_{1,n})^q)$, the q-th moment of the total mass of the correctly renormalized critical Mandelbrot cascade measure. As noted in [10], the fact that this is uniformly bounded in n for a fixed $q<1$ follows from [30, 41]. Thus $\mathbb{E}(S_n^q)$ is also uniformly bounded in n for $q<1$. So, recalling that $s_n=\sqrt{\epsilon\log n}$ and $k_n=\frac{1}{\log 2}\log n$, we conclude that for any $\epsilon\in(0,1)$, there are constants $C(\frac{\epsilon}{2})$ and $C(\epsilon)$ so that if we take n large enough, then $1-\mathbb{P}(\mathcal{B}_n)\leq C(\frac{\epsilon}{2})(2Ck_ne^{2\sqrt{2}s_n})^{1-\frac{\epsilon}{2}}n^{(1-\frac{\epsilon}{2})(\gamma-\frac{1}{2})}\leq C(\epsilon)n^{(1-\epsilon)(\gamma-\frac{1}{2})}$. Thus all we are left with is to estimate of the Laplace transform of S_n.

We make use of the following formula, valid for all non-negative random variables X:

$$1-\mathbb{E}(\exp(-\alpha X))=\int_0^\infty \alpha e^{-\alpha t}\mathbb{P}(X\geq t)dt.$$

In this formula, we set $\alpha=2Ce^{2\sqrt{2}s_n}k_nn^{(\gamma-\frac{1}{2})}$ and $X=S_n$. Recalling from the argument above that $\mathbb{E}(S_n^q)$ is uniformly bounded in n for $q<1$, by Chebyshev's inequality we see that for any $q<1$

$$\mathbb{P}(S_n\geq t)\leq C_qt^{-q}.$$

Making the change of variable $\tau=\alpha t$, we get

$$1-\mathbb{E}(e^{-\alpha S_n})\leq C_q\alpha^q\int_0^\infty e^{-\tau}\tau^{-q}d\tau.$$

Recalling again that $s_n=\sqrt{\epsilon\log n}$ and $k_n=\frac{1}{\log 2}\log n$, we see that since the integral converges, we can take q so close to one that we get

$$1-\mathbb{E}(e^{-\alpha S_n})\leq C'(\epsilon)n^{(\gamma-\frac{1}{2})(1-\epsilon)},$$

which finishes the proof of Lemma 14. □

Theorem 2 now follows quickly. We first prove the analogous statement for the measure $\nu_{\sqrt{2}}$.

Theorem 16. *For any interval $I\subset[0,1]$ and $\gamma<\frac{1}{2}$, almost surely*

$$\nu_{\sqrt{2}}(I)\leq C(\omega)\left(\log(1+|I|^{-1})\right)^{-\gamma} \tag{31}$$

where $C(\omega)$ is an almost surely finite random constant.

Proof. It is enough to restrict to dyadic subintervals. Pick $\gamma \in (0, \frac{1}{2})$. Let l be an integer so that $l(\gamma - \frac{1}{2}) < -2$. We then have, by Lemma 14, that

$$\sum_{k=1}^{\infty} \mathbb{P}\big(\max_{\sigma \in \Sigma_{k^l}^{(e/o)}} \nu_{\sqrt{2}}(I_\sigma) \geq k^{-l\gamma}\big) \leq C \sum_{k=1}^{\infty} k^{l\frac{\gamma - \frac{1}{2}}{2}} < \infty.$$

By Borel–Cantelli,

$$\max_{\sigma \in \Sigma_{k^l}^{(e/o)}} \nu_{\sqrt{2}}(I_\sigma) \leq C(\omega) k^{-l\gamma}$$

for a random (almost surely finite) constant $C(\omega)$. Combining the estimates for even and odd intervals we get

$$\max_{\sigma \in \Sigma_{k^l}} \nu_{\sqrt{2}}(I_\sigma) \leq C'(\omega) k^{-l\gamma}.$$

We note that $\max_{\sigma \in \Sigma_n} \nu_{\sqrt{2}}(I_\sigma)$ is decreasing in n so for $k^l \leq n \leq (k+1)^l$ we have

$$\max_{\sigma \in \Sigma_n} \nu_{\sqrt{2}}(I_\sigma) \leq \max_{\sigma \in \Sigma_{k^l}} \nu_{\sqrt{2}}(I_\sigma) \leq C'(\omega) k^{-l\gamma} \leq C'(\omega) 2^{l\gamma} n^{-\gamma},$$

which is the desired result. ☐

Proof of Theorem 2. As in the proof of Lemma 12 we note that

$$\mu_{\sqrt{2}}(I) \leq e^{\sqrt{2}\max_{x \in [0,1]} X_0(x) - 1} \nu_{\sqrt{2}}(I),$$

where $(X_0(x))_{x \in [0,1]}$ is a Gaussian process with a continuous covariance kernel. The quantity $e^{\sqrt{2}\max_{x \in [0,1]} X_0(x)}$ is almost surely finite, so Theorem 16 implies the result. ☐

4. ON THE $\mu_{\sqrt{2}}$-ALMOST EVERYWHERE LOCAL BEHAVIOR OF $\mu_{\sqrt{2}}$

In this section we consider the following question: what can be said of the size of smallest possible sets of full $\mu_{\sqrt{2}}$-measure? This question is partially answered by Theorem 3, which is proven in this section.

Let $f : \mathbb{N} \to \mathbb{R}^+$ be an ultimately nonincreasing function tending to 0 at infinity and consider the sets

$$E_n^f = \{x : \mu_{\sqrt{2}}(I_n(x)) \leq f(n)\}.$$

We will determine a class of functions f for which we have

$$\sum_n \mu_{\sqrt{2}}\left(E_n^f\right) < \infty \quad \text{almost surely.}$$

For a nontrivial result it is already enough to consider the expectation of the series above. We fix a sequence $(\eta_n)_{n \geq 1}$ taking values in $(0, 1)$ and write

$$\mu_{\sqrt{2}}\left(E_n^f\right) = \int_0^1 \mathbf{1}_{\{\mu_{\sqrt{2}}(I_n(x)) \leq f(n)\}} \mu_{\sqrt{2}}(\,\mathrm{d}x) = \sum_{\sigma \in \Sigma_n} \mu_{\sqrt{2}}(I_\sigma) \mathbf{1}_{\{\mu_{\sqrt{2}}(I_\sigma) \leq f(n)\}}$$

$$\leq \sum_{\sigma \in \Sigma_n} \mu_{\sqrt{2}}(I_\sigma) \left(\frac{f(n)}{\mu_{\sqrt{2}}(I_\sigma)}\right)^{\eta_n} = \sum_{\sigma \in \Sigma_n} \mu_{\sqrt{2}}(I_\sigma)^{1-\eta_n} f(n)^{\eta_n}.$$

Let $\epsilon_n = -\dfrac{\log(f(n))}{n}$ take the form $\gamma\sqrt{\dfrac{\log(n)}{n} + \alpha\dfrac{\log\log(n)}{n}}$ for $n \geq 3$, where $\alpha > 0$ and $\gamma > 0$ are to be prescribed. Assume $\eta_n = \lambda\epsilon_n$.

Denoting by W_n the n-th level lognormal factor

$$W_n \overset{d}{=} \exp\left(\sqrt{2}X_n - \mathbb{E}X_n^2\right) \overset{d}{=} \exp\left(\sigma_n N - \frac{\sigma_n^2}{2}\right),$$
$$N \sim N(0,1), \quad \sigma_n^2 = 2n\log 2,$$

we have, for each $\sigma \in \Sigma_n$, $\mu_{\sqrt{2}}(I_\sigma) \overset{d}{=} 2^{-n}W_nY_n$ where Y_n is a copy of Y independent of W_n. Moreover, by Theorem 1 we have $\mathbb{E}(Y^{1-\eta}) = O(\eta^{-1})$ as $\eta \to 0^+$. These remarks yield

$$\mathbb{E}\mu_{\sqrt{2}}\left(E_n^f\right) \leq 2^n 2^{-n(1-\eta_n)}\mathbb{E}(W_n^{1-\eta_n})\mathbb{E}(Y^{1-\eta_n})e^{-n\epsilon_n\eta_n} \leq Ce^{n(\log(2)\eta_n^2 - \epsilon_n\eta_n) - \log(\eta_n)}.$$

A computation yields for $n \geq 3$

$$n(\log(2)\eta_n^2 - \epsilon_n\eta_n) - \log(\eta_n) = \big(c + \frac{1}{2}\big)\log(n) + \big(c\alpha - \frac{1}{2}\big)\log\log(n) + O(1),$$

where $c = \log(2)\lambda^2\gamma^2 - \lambda\gamma^2$. With the order of magnitude chosen for ϵ_n, taking $c = -\frac{3}{2}$ is optimal in view of making $\sum_{n\geq 1}\mathbb{E}\mu_{\sqrt{2}}\left(E_n^f\right)$ convergent. This condition requires the equation $\log(2)\lambda^2\gamma^2 - \lambda\gamma^2 + \frac{3}{2} = 0$ to have solutions in λ. This imposes $\gamma \geq \sqrt{6\log(2)}$, hence we choose $\gamma = \sqrt{6\log(2)}$ to minimize ϵ_n. It then turns out that if $-\frac{3}{2}\alpha - \frac{1}{2} < -1$, i.e. $\alpha > \frac{1}{3}$, then $\sum_{n\geq 1}\mathbb{E}\mu_{\sqrt{2}}\left(E_n^f\right) < \infty$.

Theorem 3 follows from the preceding estimates by an application of the Borel–Cantelli lemma to the measure $\mu_{\sqrt{2}}$. As an application of Theorem 3 we present the following simple corollary.

Corollary 17. *Almost surely, there exists a set of Hausdorff dimension* 0 *that has full* $\mu_{\sqrt{2}}$*-measure.*

Proof. Let

$$E = \left\{x : \mu_{\sqrt{2}}(I_n(x)) \geq f(n) \text{ for all but finitely many } n\right\},$$

where $f = f_\alpha$ for some $\alpha > \frac{1}{3}$. Since, by Theorem 3, E almost surely has full $\mu_{\sqrt{2}}$-measure, we only need to show that a.s. it has Hausdorff dimension 0.

Let $\{I_\sigma\}_{\sigma\in\Sigma_n^f}$ be the collection of dyadic subintervals of $[0,1]$ such that $|\sigma| \geq n$ and $\mu_{\sqrt{2}}(I_\sigma) \geq f(|\sigma|)$. Clearly, for any n, $\{I_\sigma\}_{\sigma\in\Sigma_n^f}$ is a cover of E. But for any $s > 0$ and sufficiently large $n \in \mathbb{N}$ we have $2^{-\frac{s}{2}|\sigma|} \leq \mu_{\sqrt{2}}(I_\sigma)$ for all $\sigma \in \Sigma_n^f$, so

$$\begin{aligned}\sum_{\sigma\in\Sigma_n^f} |I_\sigma|^s &= \sum_{k\geq n}\sum_{\sigma\in\Sigma_n^f, |\sigma|=k} |I_\sigma|^s = \sum_{k\geq n}\sum_{\sigma\in\Sigma_n^f, |\sigma|=k} |I_\sigma|^{\frac{s}{2}}(2^{-|\sigma|})^{\frac{s}{2}} \\ &\leq \sum_{k\geq n} 2^{-k\frac{s}{2}} \sum_{\sigma\in\Sigma_n^f, |\sigma|=k} \mu_{\sqrt{2}}(I_\sigma) \\ &\leq \mu_{\sqrt{2}}([0,1]) \sum_{k\geq n} 2^{-k\frac{s}{2}}.\end{aligned}$$

The last expression tends to 0 as $n \to \infty$. It follows that for any $s > 0$ the set E has zero Hausdorff s-measure, which implies the claim. ☐

5. Appendix: Scale invariance properties

In this section we give the computations leading to the statements (2) and (6) on the exact scale invariance of the field X and of the measure $\mu_{\sqrt{2}}$. We also discuss the $\star$-scaling relation for the measure $\nu_{\sqrt{2}}$ given in (19).

Proposition 18. *The random measure $\mu_{\sqrt{2}}$ satisfies the exact scale invariance property* (6)*, i.e. for any interval $I \subset [0,1]$*

$$\mu_{\sqrt{2}}\lfloor I \overset{d}{=} |I| e^{\sqrt{2}X(I) - \mathbb{E}X(I)^2} \mu_{\sqrt{2}}^I,$$

where $\mu_{\sqrt{2}}\lfloor I$ denotes the restriction of $\mu_{\sqrt{2}}$ onto I and $\mu_{\sqrt{2}}^I$ is a random measure independent of $X(I)$ with the law given by

$$\Big(\mu_{\sqrt{2}}(A)\Big)_{A\in\mathcal{B}(I)} \overset{d}{=} \Big(\mu_{\sqrt{2}}(|I|^{-1}A)\Big)_{A\in\mathcal{B}(I)}.$$

Remark. Writing the scaling relation simultaneously for a set $\{I_j\}$ of subintervals of $[0,1]$, one has

$$\Big(\mu_{\sqrt{2}}\lfloor I_j\Big)_j \overset{d}{=} \Big(|I_j| e^{\sqrt{2}X(I_j) - \mathbb{E}X(I_j)^2} \mu_{\sqrt{2}}^{I_j}\Big)_j,$$

where the $\mu_{\sqrt{2}}^{I_j}$ are random measures such that for each j,

$$\Big(\mu_{\sqrt{2}}^{I_j}(J)\Big)_{J\in\mathcal{B}(I_j)} \overset{d}{=} \Big(\mu_{\sqrt{2}}(|I_j|^{-1}J)\Big)_{J\in\mathcal{B}(I_j)} \quad \text{and} \quad \mu_{\sqrt{2}}^{I_j} \perp \{X(A)\}_{A\subset\mathcal{C}(I_j)}.$$

However we stress that for subintervals of the unit interval, for $j \neq k$ the measure $\mu_{\sqrt{2}}^{I_j}$ is not independent either of $\mu_{\sqrt{2}}^{I_k}$ or $X(I_k)$.

Proof. We first show that (2) holds. Consider, for notational convenience, the interval $I = [0, y]$ with $0 < y < 1$. By definition, for $t \geq \log 1/y$ we have

$$(X_t(x))_{x\in I} = (X(I) + X_t^I(x))_{x\in I}.$$

Therefore it suffices to check that

$$(X_t^I(x))_{x\in I} \overset{d}{=} (X_{t-\log 1/y}(x/y))_{x\in I}$$

and since the processes are Gaussian, it is enough to consider the covariance structures. Checking that the covariances of the processes are the same is demonstrated in Figure 1.



FIGURE 1. *Left.* The sets $\mathcal{C}_t(x_1) \setminus \mathcal{C}_t(x_2)$ and $\mathcal{C}_t(x_2) \setminus \mathcal{C}_t(x_1)$ are shaded light gray, while the intersection $\big(\mathcal{C}_t(x_1) \cap \mathcal{C}_t(x_2)\big) \setminus \mathcal{C}([0,y])$ is dark gray. The law of the Gaussian process $(X_t^I(x))_{x\in[0,y]}$ is determined by the hyperbolic areas of these sets for all pairs $(x_1,x_2) \in [0,y]^2$. The set $\mathcal{C}([0,y])$, contained in every $\mathcal{C}_t(x)$ for $x \in [0,y]$, has been left white. *Right.* Closing the gap left by the set $\mathcal{C}([0,y])$ does not affect the hyperbolic areas of any of the shaded regions. Scaling this picture by $1/y$ also leaves the hyperbolic areas invariant, giving the distributional equality $(X_t^I(x))_{x\in I} \overset{d}{=} (X_{t-\log 1/y}(x/y))_{x\in I}$.

Showing the exact scale invariance of $\mu_{\sqrt{2}}$ is now simple, as one only needs to note that the measure defined analogously to the subcritical measures vanishes: for any intervals $J \subset I \subset [0,1]$ we have

$$\begin{aligned}
\mu_{\sqrt{2}}(J) &= \lim_{t\to\infty} \int_J \Big(\sqrt{2}(t+1) - X_t(x)\Big)\, e^{\sqrt{2}X_t(x) - \mathbb{E}X_t(x)^2}\,\mathrm{d}x \\
&= \lim_{t\to\infty} \int_J \Big(\sqrt{2}\mathbb{E}X(I)^2 - X(I)\Big)\, e^{\sqrt{2}X_t(x) - \mathbb{E}X_t(x)^2}\,\mathrm{d}x \\
&\quad + \lim_{t\to\infty} \int_J \Big(\sqrt{2}(t+1-\mathbb{E}X(I)^2) - X_t^I(x)\Big)\, e^{\sqrt{2}X_t(x) - \mathbb{E}X_t(x)^2}\,\mathrm{d}x \\
&= 0 \\
&\quad + e^{\sqrt{2}X(I) - \mathbb{E}X(I)^2} \lim_{t\to\infty} \int_J \Big(\sqrt{2}(t+1-\mathbb{E}X(I)^2) - X_t^I(x)\Big)\, e^{\sqrt{2}X_t^I(x) - \mathbb{E}X_t^I(x)^2}\,\mathrm{d}x \\
&=: |I| e^{\sqrt{2}X(I) - \mathbb{E}X(I)^2} \mu^I(|I|^{-1}J),
\end{aligned}$$

where μ^I a random measure with the law of μ and independent of $X(I)$. Note that the measure μ^I defined here depends on the field X only through the processes $(X_t^I(x))_{x\in I}$, $t > 0$. This observation implies the statement on the simultaneous scaling relations for a set of intervals $\{I_j\}$. □

We then consider $\star$-scale invariance, as defined in [2], and the measure $\nu_{\sqrt{2}}$ defined for the proof of Theorem 2. A random measure ν on $[0,1]$ is called $\star$-scale invariant on scale $\epsilon \in (0,1]$ if there exist a process $(\omega_\epsilon(x))_{x\in[0,1]}$ and a random measure ν^ϵ that are indepedent of each other and satisfy

$$(\nu(A))_{A\in\mathcal{B}([0,1])} \overset{d}{=} \left(\int_A e^{\omega_\epsilon(x)}\,\mathrm{d}\nu^\epsilon(x)\right)_{A\in\mathcal{B}([0,1])}$$



FIGURE 2. The cones $\mathcal{C}_t(x_1)$ and $\mathcal{C}_t(x_2)$ have been shaded gray, with the parts in $\mathcal{C}_{\log\frac{1}{\epsilon}}(x_1)$ and $\mathcal{C}_{\log\frac{1}{\epsilon}}(x_2)$ highlighted. By scaling the part of the picture below the line $\log\frac{1}{\epsilon}$ by ϵ^{-1} we get the equality of distributions

$$(Y_t(x) - Y_{\log\frac{1}{\epsilon}}(x))_{x\in[0,1]} \overset{d}{=} (Y_{t-\log\frac{1}{\epsilon}}(\epsilon^{-1}x))_{x\in[0,1]}.$$

This immediately implies (32), since the process $(Y_{\log\frac{1}{\epsilon}}(x))_{x\in[0,1]}$ is independent of $(Y_t(x) - Y_{\log\frac{1}{\epsilon}}(x))_{x\in[0,1]}$.

and

$$\left(\nu^{\epsilon}(A)\right)_{A\in\mathcal{B}([0,1])} \overset{d}{=} \left(\nu(\epsilon^{-1}A)\right)_{A\in\mathcal{B}([0,1])}.$$

The measure

$$\nu_{\sqrt{2}}(\,\mathrm{d}x) = \lim_{t\to\infty} \sqrt{t}e^{\sqrt{2}Y_t(x)-\mathbb{E}Y_t(x)^2}\,\mathrm{d}x,$$

where $Y_t(x) = X_t(x) - X_0(x) = W(\mathcal{C}_t(x)\setminus\mathcal{C}_0(x))$, is $\star$-scale invariant on every scale $\epsilon\in(0,1]$ with

$$\omega_\epsilon(x) = \sqrt{2}Y_{\log\frac{1}{\epsilon}}(x) + 2\log\epsilon.$$

This can be seen by first deducing the scale invariance property

$$(Y_t(x))_{x\in[0,1]} \overset{d}{=} \left(Y_{\log\frac{1}{\epsilon}}(x) + Y'_{t-\log\frac{1}{\epsilon}}(\epsilon^{-1}x)\right)_{x\in[0,1]}, \tag{32}$$

where Y' is an independent realization of the field Y, from Figure 2 and then performing a computation analogous to the one above for $\mu_{\sqrt{2}}$.

REFERENCES

[1] E. Aïdékon and Z. Shi: The Seneta-Heyde scaling for the branching random walk, arXiv:1102.0217.

[2] R. Allez, R. Rhodes, V. Vargas, Lognormal scale invariant random measures, To appear in Prob. Th. Relat. Fields, arXiv:1102.1895v3.

[3] K. Astala, P. Jones, A. Kupiainen, E. Saksman: Random conformal weldings, Acta Math. 207, 2, (2011), 203–254.

[4] E. Bacry and J.-F. Muzy, Log-infinitely divisible multifractal processes, Commun. Math. Phys., **236** (2003), 449–475.

[5] J. Barral and A.H. Fan, Covering numbers of different points in the Dvoretzky covering, Bull. Sci. Math., **129** (2005), 275–317.

[6] J. Barral, X. Jin: On exact scaling log-infinitely divisible cascades. arXiv:1208:2221

[7] J. Barral, X. Jin, R. Rhodes, V. Vargas: Gaussian multiplicative chaos and KPZ duality, arXiv:1202.5296, to appear in Comm. Math. Phys.

[8] J. Barral, R. Rhodes, V. Vargas: Limiting laws of supercritical branching random walks, C. R. Acad. Sci. Paris, Ser., I 350 (2012), 535–538.

[9] J. Barral and B.B. Mandelbrot, Multifractal products of cylindrical pulses, Probab. Theory Relat. Fields, **124** (2002), 409–430.

[10] J. Barral, A. Kupiainen, M. Nikula, E. Saksman, C. Webb: Critical Mandelbrot Cascades. arXiv:1206.5444

[11] I. Benjamini, O. Schramm: KPZ in One Dimensional Random Geometry of Multiplicative Cascades, Commun. Math. Phys., 289 (2009), 653–662.

[12] D. Buraczewski: On invariant measures of stochastic recursions in a critical case. Ann. Appl. Probab. 17 (2007), no. 4, 1245–1272.

[13] D. Buraczewski: On tails of fixed points of the smoothing transform in the boundary case. Stochastic Process. Appl. 119 (2009), no. 11, 3955–3961.

[14] R. Durrett, T. Liggett: Fixed points of the smoothing transformation. Z. Wahrsch. Verw. Gebiete 64 (1983), no. 3, 275-301.

[15] Duplantier B.: A rigorous perspective on Liouville quantum gravity and KPZ, in *Exact Methods in Low-dimensional Statistical Physics and Quantum Computing*, J. Jacobsen, S. Ouvry, V. Pasquier, D. Serban, and L.F. Cugliandolo, eds., Lecture Notes of the Les Houches Summer School: Volume 89, July 2008, Oxford University Press (Clarendon, Oxford) (2010).

[16] B. Duplantier, S., Sheffield, Duality and KPZ in Liouville Quantum Gravity, Phys. Rev. Let. **102**, 150603 (2009).

[17] B. Duplantier, S. Sheffield: Liouville quantum gravity and KPZ, Invent. Math., **185** (2011), 333–393.

[18] B. Duplantier, R. Rhodes, S. Sheffield, V. Vargas: Critical Gaussian Multiplicative Chaos: Convergence of the Derivative Martingale. arXiv:1206.1671

[19] B. Duplantier, R. Rhodes, S. Sheffield, V. Vargas: Renormalization of Critical Gaussian Multiplicative Chaos and KPZ formula. arXiv:1212:0529

[20] Y. Guivarch: Sur une extension de la notion de loi semi-stable, Ann. Inst. H. Poincaré Probab. Statist. 26 (1990), 261–285.

[21] A.-H. Fan, Sur le chaos de Lévy d'indice $0 < \alpha < 1$, *Ann. Sciences Math. Québec*, vol 21 no. 1, 1997, p. 53-66.

[22] A.-H. Fan, Limsup deviations on trees, Anal. Th. Appl., **20** (2004), 113–148.

[23] J.-P. Kahane, Sur le chaos multiplicatif, Ann. Sc. Math. Québec **9** (1985), 105–150.

[24] J.P. Kahane, Some Random Series of Functions. Cambridge University Press, 5, 1985.

[25] J.-P. Kahane, Positive martingales and random measures, Chi. Ann. of Math., **8B1** (1987), 1–12.

[26] J.-P. Kahane, Multiplications aléatoires et dimensions de Hausdorff. Ann. Inst. Henri Poincaré Probab. Stat., **23** (1987), 289–296.

[27] J.-P. Kahane, Recouvrement aléatoire et théorie du potentiel, Coll. Math., **LX/LXI** (1990), 387–411.

[28] J.-P. Kahane, J. Peyrière, Sur certaines martingales de B. Mandelbrot, Adv. Math., **22** (1976), 131–145.

[29] Q. Liu: On generalized multiplicative cascades. Stochastic Process. Appl. 86 (2000), no. 2, 263–286.

[30] T. Madaule: Convergence in law for the branching random walk seen from its tip. arXiv:1107.2543

[31] B.B Mandelbrot: Possible refinement of the lognormal hypothesis concerning the distribution of energy in intermittent turbulence, Statistical Models and Turbulence. In Rosenblatt, M. and Atta, C.V. ed. Lectures Notes in Physics. 12 (1972), 333–351. Springer–Verlag, New York.

[32] B.B Mandelbrot: Multiplications aléatores itérées et distributions invariantes par moyennes pondérées. C. R. Acad. Sci. Paris, 278 (1974), 289–292 and 355–358.

[33] B.B. Mandelbrot: Intermittent turbulence in self-similar cascades, divergence of high moments and dimension of the carrier, J. Fluid. Mech., 62 (1974), 331–358.
[34] B.B. Mandelbrot, Multifractal measures, especially for the geophysicists, PAGEOPH, **131** (1989), 5–42.
[35] B.B. Mandelbrot, Fractals and Scaling in Finance: Discontinuity, Concentration, Risk. Springer, 1997.
[36] S. Port, C. Stone: Potential theory of random walks on Abelian groups. Acta Math. 122 (1969), 19–114.
[37] D. Revuz, M. Yor, Continuous Martingales and Brownian Motion, Springer, third Edition, 1999.
[38] R. Rhodes, V. Vargas, KPZ formula for log-infinitely divisible multifractal random measures, ESAIM: Probab. & Stat., **15** (2011), 358–371.
[39] R. Robert, V. Vargas, Hydrodynamic Turbulence and Intermittent Random Fields, *Communications in Mathematical Physics*, **284** (3) (2008), 649-673.
[40] R. Robert, V. Vargas, Gaussian Multiplicative Chaos revisited, *Annals of Probability*, **38** 2 (2010), 605-631.
[41] C. Webb: Exact asymptotics of the freezing transitions of a logarithmically correlated random energy model. J. Stat. Phys, 145 (2011), 1595–1619.

LAGA (UMR 7539), Département de Mathématiques, Institut Galilée, Université Paris 13, 99 avenue Jean-Baptiste Clément , 93430 Villetaneuse, France
E-mail address: `barral@math.univ-paris13.fr`

University of Helsinki, Department of Mathematics and Statistics, P.O. Box 68 , FIN-00014 University of Helsinki, Finland
E-mail address: `antti.kupiainen@helsinki.fi`

University of Helsinki, Department of Mathematics and Statistics, P.O. Box 68 , FIN-00014 University of Helsinki, Finland
E-mail address: `miika.nikula@helsinki.fi`

University of Helsinki, Department of Mathematics and Statistics, P.O. Box 68 , FIN-00014 University of Helsinki, Finland
E-mail address: `eero.saksman@helsinki.fi`

University of Helsinki, Department of Mathematics and Statistics, P.O. Box 68 , FIN-00014 University of Helsinki, Finland
E-mail address: `christian.webb@helsinki.fi`